Mechanika Kwantowa dla doktorantów zestaw 5 na dzień 12.11.2013 godz. 13:45 sala 431a

- 1. (Complete the problem fom Nov. 5) Find a classical action for the harmonic oscillator with an external force F(t). Take limit $\omega \to 0$ to obtain classical action for a particle moving in an external force. Finally take limit $F \to 0$ to obtain an action of a free particle.
- 2. For a classical action derived in 1. take constant force F(t) = F and calculate all integrals over time. For a constant force one can derive an action in a different way by observing that the lagrangian can be rewritten in a new variable y as a free oscillator plus a constant. Variable y corresponds to a suitably shifted variable x. Derive an action in that way and compare with the previous result.
- 3. One slit diffraction.

Consider one dimensional problem of a particle moving from x = 0 at t = 0 to point (x, t) through a screen which the particle reaches at time $t_1 < t$. The screen has a hole of size 2b centered around x_0 . (This process can be understood in the following way: at time t_1 we block all possible trajectories except those passing through a hole). Calculate the propagator for such a motion.

In order to simplify calculactions replace the sharp hole by a "Gaussian hole"

$$G(x) = \begin{cases} 0 & \text{for } x < x_0 - b \text{ or } x_0 + b < x \\ & \to G(x) = e^{-(x - x_0)^2/2b^2}. \\ 1 & \text{for } x_0 - b \le x \le x_0 + b \end{cases}$$

What is the width of the image of this "Gaussian hole" at time t? How does it compare with the classical expectation?

Since the propagation before and after the slit is free, we can use the formula for a free particle propagator

$$K_0(b,a) = \sqrt{\frac{m}{2\pi i\hbar(t_b - t_a)}} e^{\frac{i}{\hbar}\frac{m}{2}\frac{(x_b - x_a)^2}{t_b - t_a}}$$

and the formula for "folding" the propagators

$$K(b,a) = \int dx_c K(b,c) K(c,a).$$

Here $a = (x_a, t_a)$ etc.

This problem can be found in the book of Feynman and Hibbs "Quantum Mechanics and Path Integrals" chapt. 3.2.