

Mechanika Kwantowa dla doktorantów  
zestaw 11 – 5.1.2017 at 8:15

1. *Instanton determinant.* In this problem we will calculate explicitly the ratio of determinants

$$K' = \frac{\det' \left( -\frac{d^2}{d\tau^2} + V''(\bar{x}(\tau)) \right)}{\det \left( -\frac{d^2}{d\tau^2} + 1 \right)},$$

where a double well potential  $V(x)$  reads:  $V(x) = \kappa(a^2 - x^2)^2$  with  $\kappa = 1/(8a^2)$ . Prime at the determinant means that the zero eigenvalue (zero mode) is not included, by  $\bar{x}(\tau)$  we denote classical trajectory.

- The eigenequation for quantum fluctuations around the classical trajectory (with  $\tau_1 = 0$ , where  $\tau_1$  is the time when the classical trajectory passes through zero):

$$\left[ -\frac{d^2}{d\tau^2} + V''(\bar{x}(\tau)) \right] y_n(\tau) = \lambda_n y_n(\tau) \quad (1)$$

corresponds to the Schrödinger equation for a potential  $U(\tau) = -3/(2\cosh^2(\tau/2))$  (where  $\tau$  plays a role of a spacial variable) and energy  $E_n = \lambda_n - 1$ , which is discussed in the "Quantum Mechanics" of Landau and Lifschitz (probl. 5 page. 81 and probl. 4 page 88, Polish edition PWN 1979).

Transform equation (1) into a hypergeometric equation for function  $w_n$  defined below:

$$y_n(\tau) = e^{\alpha\tau} w_n(\tau),$$

where

$$\alpha = \pm\sqrt{-E_n}, \quad E_n = \lambda_n - 1.$$

Show that the solution reads:

$$y_n(\tau) = \mathcal{N} \left( 3 \tanh^2 \left( \frac{\tau}{2} \right) - 6\alpha \tanh \left( \frac{\tau}{2} \right) + (4\alpha^2 - 1) \right) e^{\alpha\tau}.$$

To this end introduce the following new variables:

$$\begin{aligned} z &= \tanh \left( \frac{\tau}{2} \right), \\ u &= \frac{1}{2}(1 + z). \end{aligned}$$

In terms of variable  $u$  Eq.(1) corresponds to the hypergeometric equation

$$u(1-u)w''(u) + \{c - (a+b+1)u\}w'(u) - abw(u) = 0,$$

whose solutions are given in terms of a series

$$w(u) = F(a, b, c; u) = 1 + \frac{ab}{c} \frac{u}{1!} + \frac{a(a+1)b(b+1)}{c(c+1)} \frac{u^2}{2!} + \dots$$

- Find discrete spectrum of the bound states ( $E < 0$ ) for (1). Conditions that solutions vanish at  $\tau = \pm\infty$  give quantization of  $\alpha$ .
- **This part will be most probably shifted to next class.** To find contribution from the continuous spectrum we show first that there is no reflection for the particles scattering over the potential  $U(\tau)$ . To this end find asymptotics for two types of the solutions:  $\alpha = ik$  and  $\alpha = -ik$  in the limit  $\tau \rightarrow \pm\infty$ .
- If there is no reflection then the wave function  $y_k(\tau)$  that asymptotically behaves as  $e^{ik\tau}$  for  $\tau \rightarrow \infty$ , in the limit of  $\tau \rightarrow -\infty$  behaves as  $e^{ik\tau+i\delta_k}$ , where  $\delta_k$  is a phase shift. Show that

$$e^{i\delta_k} = \frac{1+ik}{1-ik} \frac{1+2ik}{1-2ik}.$$

Identical argument applies to the wave function, that asymptotically behaves as  $e^{-ik\tau}$ .

- Close the system in a box  $-T/2 < \tau < T/2$ . Then the wave function inside the box is a superposition of two linearly independent solutions

$$y_n(\tau) = Ay_{\alpha=ik}(\tau) + By_{\alpha=-ik}(\tau)$$

which vanishes at the boundaries

$$y_n(\pm T/2) = 0. \quad (2)$$

If the box is large, it is enough to use asymptotic forms of  $y_{\alpha=\pm ik}(\tau)$ . Show that condition (2) leads to

$$Tk - \delta_k = \pi n.$$

Let's denote solution to this equation by  $\tilde{k}_n$ . Similarly, for the Euclidean harmonic oscillator analogous solutions read  $k_n = \pi n/T$ .

- The contribution to  $K'$  coming from the continuous spectrum,  $K_{cont}$ , reads:

$$K_{cont} = \frac{\prod \tilde{\lambda}_n}{\prod \lambda_n} = \prod_{n=1}^{\infty} \frac{1 + \tilde{k}_n^2}{1 + k_n^2} = \exp\left(\sum_n \ln \frac{1 + \tilde{k}_n^2}{1 + k_n^2}\right) \approx \exp\left(\sum_n \frac{2k_n(\tilde{k}_n - k_n)}{1 + k_n^2}\right).$$

- To calculate last sum under exponent go to the continuum limit  $T \rightarrow \infty$  and convert the sum into the integral:

$$\dots = \exp\left(\frac{1}{\pi} \int_0^{\infty} dk \frac{2\delta_k k}{1 + k^2}\right) = \frac{1}{9}.$$

Last equality can be obtained by integration by parts and the explicit form of  $\delta_k$ . Full result for  $K'$  is obtained by multiplying  $K_{cont}$  by a non-zero  $\lambda$  value from the discrete part.

- Literature:

S. Coleman, *Aspects of Symmetry*, Cambridge University Press (1988), Section 7, Appendix 1.

A.I. Vainshtein, V.I. Zakharov, V.A. Novikov and M.A. Shifman, *ABC of Instantons*, Sov. Phys. Usp. **24**, 195 (1982) [Usp. Fiz. Nauk **136**, 553 (1982)].

T. Schafer and E.V. Shuryak, *Instantons in QCD*, Rev. Mod. Phys. **70** (1998) 323 [arXiv:hep-ph/9610451].