



Beyond Minimal Flavour Violation

Andrzej J. Buras
(Technical University Munich, TUM-IAS)

Zakopane Lectures 2010

Lecture I

- 1.** Grand View
- 2.** TH Framework

Lecture II

- 3.** Minimal Flavour Violation
- 4.** Motivations for BSM and BMFV
- 5.** Models for BMFV
(SUSY, LHT, RS, 4G)

Lecture III

- 6.** Patterns of Flavour Violations
BSM
- 7.** Outlook

1.

Grand View

The Standard Model

Quarks

$$\begin{pmatrix} u \\ d \end{pmatrix}_L \begin{pmatrix} c \\ s \end{pmatrix}_L \begin{pmatrix} t \\ b \end{pmatrix}_L \quad \begin{matrix} u_R & c_R & t_R \\ d_R & s_R & b_R \end{matrix} \quad \begin{matrix} + 2/3 \\ - 1/3 \end{matrix}$$

+ Leptons

Fundamental Forces

$$\left. \begin{array}{c} \text{Gauge} \\ \text{Theory} \end{array} \right\} : \underbrace{\text{SU}(3) \otimes \text{SU}(2)}_{\substack{\text{QCD} \\ \text{Strong} \\ \text{Interactions}}} \underbrace{\otimes \text{U}(1)_Y}_{\substack{\text{U}(1)_{\text{QED}} \\ \text{Electroweak Interactions}}} \quad \begin{array}{c} \text{Neutral} \\ \text{Higgs} \end{array}$$


(Gluons) (W^\pm, Z^0, γ)

Mesons

$$\begin{aligned} K^0 &= (d\bar{s}) & K^+ &= (u\bar{s}) & K^- &= (\bar{u}s) \\ \pi^+ &= (u\bar{d}) & \pi^0 &= (\bar{u}u - \bar{d}d)/\sqrt{2} & \pi^- &= (\bar{u}d) \\ B_d^0 &= (d\bar{b}) & \bar{B}_d^0 &= (\bar{d}b) & B^+ &= (u\bar{b}) \\ B_s^0 &= (s\bar{b}) & \bar{B}_s^0 &= (\bar{s}b) & B^- &= (\bar{u}\bar{b}) \end{aligned} \quad \left. \right\} \quad \begin{array}{c} q\bar{q} \\ \text{Bound} \\ \text{States} \end{array}$$

Four Basic Properties in the SM

1. Charged Current Interactions only between left-handed Quarks



$$\frac{g_2}{2\sqrt{2}} \gamma_\mu (1 - \gamma_5) \cdot V_{td}$$

2. Quark Mixing

$$\{ \text{Weak Eigenstates} \} \neq \{ \text{Mass Eigenstates} \}$$

$$\begin{bmatrix} d' \\ s' \\ b' \end{bmatrix} = \begin{bmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{bmatrix} \begin{bmatrix} d \\ s \\ b \end{bmatrix}$$



$$\begin{pmatrix} \text{Weak} \\ \text{Eigenstates} \end{pmatrix} \begin{pmatrix} \text{Unitarity} \\ \text{CKM-Matrix} \end{pmatrix} \begin{pmatrix} \text{Mass} \\ \text{Eigenstates} \end{pmatrix}$$

3. GIM Mechanism

Natural suppression of FCNC

$$\left\{ \gamma, G, Z^0, H^0 \text{ exchange between } i \text{ and } j = 0 \right\} \rightarrow \left\{ \text{Loop Induced Decays, sensitive to short distance flavour dynamics} \right\}$$

4. Asymptotic Freedom



Q

$$g_{\text{QCD}} T_{ij}^a \gamma_\mu$$

$$\alpha_{\text{QCD}} = \frac{g_{\text{QCD}}^2}{4\pi}$$

$$\alpha_{\text{QCD}}(Q) = \frac{4\pi}{\beta_0 \ln(Q^2 / \Lambda_{\overline{\text{MS}}}^2)} \left[1 - \frac{\beta_1}{\beta_0^2} \frac{\ln \ln(Q^2 / \Lambda_{\overline{\text{MS}}}^2)}{\ln(Q^2 / \Lambda_{\overline{\text{MS}}}^2)} + \dots \right]$$

$$\Lambda_{\overline{\text{MS}}}^{(5)} = 240 \pm 15 \text{ MeV} \quad \alpha_{\overline{\text{MS}}}^{(5)}(M_Z) = 0.1187 \pm 0.0009$$

SD = Short Distances (Perturbation Theory)



RG = Renormalization Group Effects



LD = Long Distances (Non-Perturbative Physics)



CP

Kobayashi-Maskawa Picture of CP Violation

CP Violation arises from **a single phase δ**
in W^\pm interactions of Quarks

ud	$c_{12}c_{13}$	us	$s_{12}c_{13}$	ub	$s_{13}e^{-i\delta}$
cd	$-s_{12}c_{23}-c_{12}s_{23}s_{13}e^{i\delta}$	cs	$c_{12}c_{23}-s_{12}s_{23}s_{13}e^{i\delta}$	cb	$s_{23}c_{13}$
td	$s_{12}s_{23}-c_{12}c_{23}s_{13}e^{i\delta}$	ts	$-s_{23}c_{12}-s_{12}s_{23}s_{13}e^{i\delta}$	tb	$c_{23}c_{13}$

Four Parameters: $(\theta_{12} \approx \theta_{\text{cabibbo}})$

$$s_{12} = |V_{us}|, \quad s_{13} = |V_{ub}|, \quad s_{23} = |V_{cb}|, \quad \delta$$

$$c_{ij} \equiv \cos \theta_{ij}; \quad s_{ij} \equiv \sin \theta_{ij}; \quad c_{13} \equiv c_{23} \equiv 1$$

Wolfenstein Parametrization

Parameters:

λ, A, ρ, η

	d	s	b
u	$1 - \frac{\lambda^2}{2}$	λ	V_{ub}
c	$-\lambda$	$1 - \frac{\lambda^2}{2}$	V_{cb}
t	V_{td}	V_{ts}	1

$$\lambda = 0.22$$

$$V_{us} = \lambda + O(\lambda^7)$$

$$V_{cb} = A\lambda^2 + O(\lambda^8)$$

$$V_{ts} = -A\lambda^2 + O(\lambda^4)$$

$$(A = 0.83 \pm 0.02)$$

$$V_{ub} \equiv A \lambda^3 (\rho - i \eta) \quad V_{td} = A \lambda^3 (1 - \bar{\rho} - i \bar{\eta})$$

$$\bar{\rho} = \rho \left(1 - \frac{\lambda^2}{2} \right)$$

$$\bar{\eta} = \eta \left(1 - \frac{\lambda^2}{2} \right)$$

(AJB, Lautenbacher, Ostermaier, 94)


$$R_b \equiv \sqrt{\bar{\rho}^2 + \bar{\eta}^2} = \left(1 - \frac{\lambda^2}{2} \right) \frac{1}{\lambda} \left| \frac{V_{ub}}{V_{cb}} \right|$$

Circle around
 $(\bar{\rho}, \bar{\eta}) = (0,0)$

$$R_t \equiv \sqrt{(1 - \bar{\rho})^2 + \bar{\eta}^2} = \frac{1}{\lambda} \left| \frac{V_{td}}{V_{cb}} \right|$$

Circle around
 $(\bar{\rho}, \bar{\eta}) = (1,0)$

Unitarity Triangle




An Important Target of Particle Physics


$$J_{CP} = \lambda^2 |V_{cb}|^2 \bar{\eta} = 2 \cdot \text{Area of unrescaled UT}$$

Area of unrescaled
UT

Tree Level Decays



Loop Induced FCNC Processes






~~CP~~ ε_K -Parameter
 $\Delta M (K_L - K_S)$

$B_d^0 - \bar{B}_d^0$ Mixing



ε'




Discovered
in 2006
(CDF, D0)








Loop Induced FCNC Processes



$$K^+ \rightarrow \pi^+ \nu \bar{\nu}, \quad K_L \rightarrow \pi^0 \nu \bar{\nu}$$




$$K_L \rightarrow \mu \bar{\mu}, \quad B_{d,s} \rightarrow \mu \bar{\mu}, \quad B \rightarrow X_S \nu \bar{\nu}$$



$$K_L \rightarrow \pi^0 e^+ e^-$$

$$B \rightarrow X_S e^+ e^-, X_S \mu \bar{\mu}$$



$$B \rightarrow X_S \gamma \quad B \rightarrow K^* \gamma$$


$$B \rightarrow X_d \gamma$$

$$b \rightarrow s \text{ gluon}$$



Hunting Δ with Rare and ~~CP~~ Decays

2011:



Transparency shown in 2003

★ Quark Mixing and CP Violation closely related in the St. Model

A diagram illustrating the relationship between CP Asymmetries in B-Decays and the parameters α , β , and γ . On the left, a green star is followed by a brace grouping "CP Asymmetries" and "in B-Decays". An arrow points from this group to a brace on the right containing α , β , and γ .

CKM Parameters from Tree-Level Decays (subject to very small NP Pollution)

$$|V_{us}| = s_{12} = 0.2254 \pm 0.0008 \quad |V_{ub}| = s_{13} = (3.9 \pm 0.4) \cdot 10^{-3}$$

$$|V_{cb}| = s_{23} = (41.2 \pm 1.1) \cdot 10^{-3} \quad \delta_{\text{CKM}} = \gamma_{\text{UT}} = (75 \pm 15)^{\circ}$$



(-phase of V_{ub})

$$(\sin 2\beta)_{\psi K_s} = 0.672 \pm 0.023$$

↑
(-phase of V_{td})

$$\rightarrow \beta = (21.1 \pm 0.9)^{\circ}$$

but could be subject to
NP pollution

$$\text{Phase of } V_{ts}: \approx - (1.2 \pm 0.1)^{\circ}$$

Hierarchical Structure of the CKM Matrix

$$\begin{pmatrix} 0.97 & s_{12} & s_{13} e^{-iy} \\ -s_{12} & 0.97 & s_{23} \\ s_{12}s_{23} - s_{13}e^{iy} & -s_{23} & 1 \end{pmatrix}$$

$$s_{13} \ll s_{23} \ll s_{12}$$

$$(4 \cdot 10^{-3}) \quad (4 \cdot 10^{-2}) \quad (0.2)$$



GIM Structure of FCNC's

Large \mathcal{CP} effects in B_d
 Small \mathcal{CP} effects in B_s
 Tiny \mathcal{CP} effects in K_L

$$A_{CP}(B_d \rightarrow \psi K_s) \approx 0(1) \quad S_{\psi K_s} \approx \frac{2}{3}$$

$$A_{CP}(B_s \rightarrow \psi \phi) \approx 0(10^{-2}) \quad S_{\psi \phi} \approx \frac{1}{25}$$

$$\varepsilon \approx 0(10^{-3}) \quad \varepsilon' \approx 0(10^{-6})$$



$$Br(K_L \rightarrow \pi^0 \bar{v} v) \approx 0(10^{-11})$$

PMNS: Negligible LFV

(tiny v masses)

Unitarity Triangle Fits

(Icons of Flavour Physics)



Impressive Success of the CKM Picture of Flavour Changing Interactions

(GIM)
(NFC)

(Once quark masses determined : only 4 parameters)

- 1.** All leading decays of K , D , B_s^0 , B_d^0 mesons correctly described
- 2.** Suppressed transitions : $K^0 - \bar{K}^0$, $B_d^0 - \bar{B}_d^0$, $B_s^0 - \bar{B}_s^0$ mixings found at suppressed level
- 3.** CP-violating Data (K , B_d) correctly described
- 4.** $B \rightarrow X_s \gamma$, $B \rightarrow X_s l^+ l^-$ OK

CP in B_s ?

(g-2) $_\mu$?

5.

Very very highly suppressed transitions in the SM
consistent with experiment: (not seen)

Standard Model	Exp Upper Bound
$\text{Br}(B_s \rightarrow \mu^+ \mu^-) \cong 3 \cdot 10^{-9}$	$\sim 4 \cdot 10^{-8}$
$\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu}) \cong 3 \cdot 10^{-11}$	$\sim 6 \cdot 10^{-8}$
$\text{Br}(K_L \rightarrow \mu e) \cong 10^{-40}$	$\sim 10^{-12}$
$\text{Br}(\mu \rightarrow e \gamma) \approx 10^{-54}$	$\sim 10^{-11}$
$d_n \approx 10^{-32} \text{ ecm}$	$\sim 10^{-26} \text{ ecm}$

Yet

Impressive Success of the CKM Picture of Flavour Changing Interactions

(GIM)

- 1.** EW-Symmetry Breaking has to be better understood.
- 2.** Hierarchies in Fermion Masses and Mixing Angles have to be understood with the help of some New Physics (NP). This NP could have impact on Low Energies.
- 3.** There is still a lot of room for NP contributions, in particular in rare decays of mesons and leptons, in \mathcal{CP} flavour violating transitions and EDM's.
- 4.** Matter-Antimatter Asymmetry \rightarrow New CP Phases needed.
- 5.** Several tensions between the flavour data and the SM exist.

Superstars of 2010 – 2015 (Flavour Physics)

$$S_{\psi\phi} \\ (B_s \rightarrow \phi\phi)$$

$$B_s \rightarrow \mu^+ \mu^- \\ (B_d \rightarrow \mu^+ \mu^-) \\ (B^+ \rightarrow \tau^+ \nu_\tau)$$

$$K^+ \rightarrow \pi^+ \bar{\nu}\bar{\nu} \\ (K_L \rightarrow \pi^0 \bar{\nu}\bar{\nu}) \\ (B_d \rightarrow K^* \mu^+ \mu^-)$$

γ
from Tree
Level
Decays

$$\mu \rightarrow e\gamma \\ \tau \rightarrow \mu\gamma \\ \tau \rightarrow e\gamma \\ \mu \rightarrow 3e \\ \tau \rightarrow 3 \text{ leptons}$$

$$\varepsilon'/\varepsilon \\ (\text{Lattice}) \\ \text{EDM's} \\ (g-2)_\mu$$

Standard Model Predictions for Superstars

$$S_{\psi\phi} = 0.035 \pm 0.005$$

$$(S_{\psi\phi})_{\text{exp}} = 0.52 \pm 0.20$$

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) = (3.2 \pm 0.3) \cdot 10^{-9}$$

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-)_{\text{exp}} \leq 4.2 \cdot 10^{-8}$$

$$\text{Br}(B_d \rightarrow \mu^+ \mu^-) = (1.0 \pm 0.1) \cdot 10^{-10}$$

$$\text{Br}(B_d \rightarrow \mu^+ \mu^-)_{\text{exp}} \leq 1.0 \cdot 10^{-8}$$

$$\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu}) = (8.4 \pm 0.7) \cdot 10^{-11}$$

$$\text{Br}(K^+ \rightarrow \pi^+ \nu \bar{\nu})_{\text{exp}} = (17 \pm 11) \cdot 10^{-11}$$

$$\gamma = (68 \pm 7)^\circ$$

$$\gamma_{\text{exp}} = (75 \pm 15)^\circ$$

$$\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu}) = (2.8 \pm 0.6) \cdot 10^{-11}$$


$$\text{Br}(K_L \rightarrow \pi^0 \nu \bar{\nu})_{\text{exp}} \leq 6 \cdot 10^{-8}$$

2.


Theoretical Framework

The Problem of Strong Interactions

$B_d^0 - \bar{B}_d^0$ Mixing (SM)



$B_d^0 - \bar{B}_d^0$ Mixing (MSSM)



SD

: Perturbative
(Asymptotic Freedom)


LD

: Non-Perturbative
(Confinement)

Effective Field Theory

Full Theory

$(W^\pm, Z^0, G, \gamma, t, H^0, b, u, d, s, c, l)$






$\mu \geq M_W$



Effective Theory


$(G, \gamma, b, u, d, s, c, l)$



$\mu \equiv 0(m_b)$


"Generalized Fermi Theory" with calculable
"couplings" $C_B(\mu), C_2(\mu), \dots$

Operator Product Expansion



$$\langle \bar{K}^0 | (\bar{s}d)_{V-A} (\bar{s}d)_{V-A} | K^0 \rangle = \frac{8}{3} \hat{B}_K F_K^2 m_K^2 [\alpha_s(\mu)]^{2/9}$$

Master Formula for FCNC Amplitudes



Possible Dirac Structures in $K^0 - \bar{K}^0$ and $B_{d,s}^0 - \bar{B}_{d,s}^0$

SM:

$$\gamma_\mu (1 - \gamma_5) \otimes \gamma^\mu (1 - \gamma_5)$$

Beyond SM:

$$\begin{aligned} & \gamma_\mu (1 - \gamma_5) \otimes \gamma^\mu (1 + \gamma_5) \\ & (1 - \gamma_5) \otimes (1 + \gamma_5) \\ & (1 - \gamma_5) \otimes (1 - \gamma_5) \\ & \sigma_{\mu\nu} (1 - \gamma_5) \otimes \sigma^{\mu\nu} (1 - \gamma_5) \end{aligned}$$

MSSM with large $\tan\beta$

General Supersymmetric Models

Models with complicated Higgs System


Warped Extra Dimensions

NLO $\left[\eta_{\text{QCD}}^i \right]^{\text{New}}$: Ciuchini, Franco, Lubicz,
Martinelli, Scimemi, Silvestrini
AJB, Misiak, Urban, Jäger

Basic Diagrams in FCNC Processes

Penguin Family


↑
(GIM broken
at one loop)



New Physics
enters here




Similar
diagrams
in LFV
and EDM's

Box
Diagrams



Tree
Diagrams

(GIM broken
at tree level)




RS

Generated
through
mixing with
New Gauge
Bosons

Double Higgs Penguin
in SUSY

Most popular BSM Directions



New gauge bosons, fermions, scalars in loops
and even trees with often non-CKM interactions.

2 x 2 Flavour Matrix of Basic NP Scenarios

(AJB, hep-ph/0101336, Erice)


	SM Operators	+ Additional Operators
CKM	A CMFV (Y_t) SM, 2 HDM at low $\tan\beta$ LH without T-parity Universal flat ED	B MFV (Y_t, Y_b) MSSM with MFV 2 HDM at large $\tan\beta$
New Flavour (CP) Violating Interactions	C beyond CMFV LH with T-parity Some Z'-models 4 th generation	D beyond MFV MSSM with $(\delta_{ij})_{AB} \neq 0$ RS, Other Z'-models, LR Models, NMfv

Little Hierarchy Problem

Electroweak Precision Tests

+

Agreement of the CKM Picture of Flavour and CP Violation with existing Data (FCNC)



(generic)

$\Lambda_{\text{NP}} \approx 5 \text{ TeV}$

Very strong Constraints on Physics beyond SM with scales $O(1 \text{ TeV})$

Necessary to solve the hierarchy problem

(generic)

$\Lambda_{\text{NP}} \approx 1000 \text{ TeV}$

$(M_{\text{PLANCK}} \gg \Lambda_{\text{EW}})$

Message 1

: **New Physics at TeV-Scale must have
a non-Generic Flavour Structure**

Message 2

: **Protection Mechanisms to
suppress FCNCs generated
by TeV-Scale New Physics
required**

Ciuchini et al
Isidori et al
Agashe et al
+50



MFV

GIM

RS-GIM

T-Parity

R-Parity

Alignment


Degeneracy


Flavour Symmetries
(abelian, non-abelian)

Custodial Symmetries

(continuous, discrete)

In our search for a more
fundamental theory we need
to improve our understanding
of Flavour





Basic Questions for Flavour Physics

New Flavour
violating
CPV phases?

Flavour Conserving
CPV phases?

Non-MFV
Interactions?

Right-Handed
Charged
Currents?

Scalars H^0, H^\pm
and related
FCNC's?

New Fermions?
New Gauge
Bosons?



How to explain dynamically 22 free
Parameters in the Flavour Sector ?

3.

Minimal Flavour Violation

General Structure in Models with Constraint Minimal Flavour Violation

Ciuchini, Degrassi, Gambino, Giudice;
AJB, Gambino, Gorbahn, Jäger, Silvestrini;

- ★ **No new Operators** (Dirac and Colour Structures) beyond those present in the SM
- ★ Flavour Changing Transitions governed by CKM. **No new complex phases** beyond those present in the SM



$$A(\text{Decay}) = B_i \eta_{\text{QCD}}^i V_{\text{CKM}}^i \left[F_{\text{SM}}^i + \underbrace{F_{\text{New}}^i}_{\text{real}} \right]$$

Minimal Flavour Violation (MFV)

MFV

SM Yukawa Couplings are the only breaking sources of the $SU(3)^5$ flavour symmetry of the low-energy effective theory

(Y_t, Y_b)

D'Ambrosio, Guidice, Isidori, Strumia (02) Chivukula, Georgi (87)



CKM the only source of Flavour Violation
but for $Y_t \approx Y_b$ new operators could enter

CMFV

Operator structure of
SM remains



AJB, Gambino, Gorbahn, Jäger, Silvestrini (00)
Ali, London

VERY STRONG
RELATIONS
BETWEEN
K and B Physics
and generally
 $\Delta F=2$ and $\Delta F=1$
FCNC Processes

Related
Studies

- : Ratz et al (08)
- : Smith et al (08)
- Zupan et al (09)
- Kagan et al (09)

Spurion Technology
Nir et al.
AGIS
Feldmann, Mannel

also beyond
MFV

Model independent Relations:

$$\frac{Br(B_s \rightarrow \mu^+ \mu^-)}{Br(B_d \rightarrow \mu^+ \mu^-)} = \frac{\hat{B}_{B_d}}{\hat{B}_{B_s}} \frac{\tau(B_s)}{\tau(B_d)} \frac{\Delta M_s}{\Delta M_d}$$

(CMFV)

$$\frac{Br(B \rightarrow X_s \nu \bar{\nu})}{Br(B \rightarrow X_d \nu \bar{\nu})} = \frac{|V_{ts}|^2}{|V_{td}|^2} = \frac{m_{B_d}}{m_{B_s}} \frac{1}{\xi^2} \frac{\Delta M_s}{\Delta M_d}$$

(CMFV)

$$(\sin 2\beta)_{B \rightarrow \psi K_S} = (\sin 2\beta)_{K \rightarrow \pi \nu \bar{\nu}}$$

(MFV)

The violation of these model independent MFV (CMFV) relations would signal new flavour and CP-violating interactions (and/or new operators)

Relations between $\Delta M_{s,d}$ and $B_{s,d} \rightarrow \mu\bar{\mu}$ in Models with Minimal Flavour Violation

(AJB, hep-ph/0303060)

$$\Delta M_q \sim \hat{B}_q F_{B_q}^2 |V_{tq}|^2 S(x_t, x_{new})$$

$$Br(B_q \rightarrow \mu\bar{\mu}) \sim F_{B_q}^2 |V_{tq}|^2 Y^2(x_t, \bar{x}_{new})$$

Large hadronic
uncertainties
due to $F_{B_q}^2$

$$F_{B_d} \sqrt{\hat{B}_d} = \begin{pmatrix} 235 \pm 33 & +0 \\ & -24 \end{pmatrix} \text{MeV} \quad F_{B_d} = (189 \pm 27) \text{ MeV}$$

$$F_{B_s} \sqrt{\hat{B}_d} = (276 \pm 38) \text{ MeV} \quad F_{B_s} = (230 \pm 30) \text{ MeV}$$

2003

$$\hat{B}_d = 1.34 \pm 0.12$$

$$\hat{B}_s = 1.34 \pm 0.12$$

$$\frac{\hat{B}_s}{\hat{B}_d} = 1.00 \pm 0.03$$

(No problems with
chiral logs and
quenching)

Relations between $\Delta M_{s,d}$ and $B_{s,d} \rightarrow \mu\bar{\mu}$ in Models with Minimal Flavour Violation

(AJB, hep-ph/0303060)

$$\Delta M_q \sim \hat{B}_q F_{B_q}^2 |V_{tq}|^2 S(x_t, x_{new})$$

$$Br(B_q \rightarrow \mu\bar{\mu}) \sim F_{B_q}^2 |V_{tq}|^2 Y^2(x_t, \bar{x}_{new})$$

Moderate hadronic
uncertainties
due to $F_{B_q}^2$

$$F_{B_d} \sqrt{\hat{B}_d} = \begin{pmatrix} 216 \pm 15 \end{pmatrix} \text{MeV} \quad F_{B_d} = (193 \pm 10) \text{ MeV}$$

$$F_{B_s} \sqrt{\hat{B}_d} = (275 \pm 13) \text{ MeV} \quad F_{B_s} = (239 \pm 10) \text{ MeV}$$

2010

$$\hat{B}_d = 1.26 \pm 0.11$$

$$\hat{B}_s = 1.33 \pm 0.06$$

$$\frac{\hat{B}_s}{\hat{B}_d} = 0.95 \pm 0.03$$

(No problems with
chiral logs and
quenching)

$\text{Br}(B_{s,d} \rightarrow \mu\bar{\mu})$ from $\Delta M_{s,d}$

$$\text{Br}(B_q \rightarrow \mu\bar{\mu}) = 4.39 \cdot 10^{-10} \frac{\tau(B_q)}{\hat{B}_q} \frac{Y^2(x_t, \bar{x}_{\text{new}})}{S(x_t, x_{\text{new}})} \Delta M_q$$

No dependence
on $F_{B_q}^2$

SM:

$$\text{Br}(B_s \rightarrow \mu\bar{\mu}) = 3.2 \cdot 10^{-9} \left[\frac{\tau(B_s)}{1.43 \text{ ps}} \right] \left[\frac{1.33}{\hat{B}_s} \right] \left[\frac{\bar{m}_t(m_t)}{164 \text{ GeV}} \right]^{1.6} \left[\frac{\Delta M_s}{17.8 / \text{ps}} \right]$$

$$\text{Br}(B_d \rightarrow \mu\bar{\mu}) = 1.0 \cdot 10^{-10} \left[\frac{\tau(B_d)}{1.52 \text{ ps}} \right] \left[\frac{1.26}{\hat{B}_d} \right] \left[\frac{\bar{m}_t(m_t)}{164 \text{ GeV}} \right]^{1.6} \left[\frac{\Delta M_d}{0.51 / \text{ps}} \right]$$

(Example)

$$\Delta M_s = (17.8 \pm 0.1 / \text{ps})$$



$$\text{Br}(B_s \rightarrow \mu\bar{\mu}) = (3.2 \pm 0.2) \cdot 10^{-9}$$

$$\Delta M_d = (0.507 \pm 0.006 / \text{ps})$$



$$\text{Br}(B_d \rightarrow \mu\bar{\mu}) = (1.0 \pm 0.1) \cdot 10^{-10}$$

Moreover new Physics Effects can be easier seen



Testing MFV through $B_{s,d} \rightarrow \mu\bar{\mu}$ and $\Delta M_{s,d}$

$$\frac{Br(B_s \rightarrow \mu\bar{\mu})}{Br(B_d \rightarrow \mu\bar{\mu})} = \frac{\hat{B}_d}{\hat{B}_s} \underbrace{\frac{\tau(B_s)}{\tau(B_d)}}_{(0.95 \pm 0.03)} \underbrace{\frac{\Delta M_s}{\Delta M_d}}_{\text{Experiment}}$$

Valid in MFV models in which only SM operators relevant.

Violation of this relation would indicate the presence of new operators and generally of non-minimal flavour violation.

4.

Motivations for BSM and BMFV

Yet

Impressive Success of the CKM Picture of Flavour Changing Interactions

(GIM)

- 1.** EW-Symmetry Breaking has to be better understood.
- 2.** Hierarchies in Fermion Masses and Mixing Angles have to be understood with the help of some New Physics (NP). This NP could have impact on Low Energies.
- 3.** There is still a lot of room for NP contributions, in particular in rare decays of mesons and leptons, in \mathcal{CP} flavour violating transitions and EDM's.
- 4.** Matter-Antimatter Asymmetry \rightarrow New CP Phases needed.
- 5.** Several tensions between the flavour data and the SM exist.

**Can SM describe simultaneously
 \cancel{CP} in K and B_d Systems?**

$$R_t^2 \approx \xi \frac{\Delta M_d}{\Delta M_s}$$

Can SM describe simultaneously \mathcal{CP} in K and B_d Systems?

$$|\varepsilon_K|^{\text{SM}} \sim \kappa_\varepsilon \hat{B}_K |V_{cb}|^2 \left(\frac{1}{2} |V_{cb}|^2 R_t^2 \underbrace{\sin 2\beta}_{\text{(NLO)}} \eta_{tt}^{\text{QCD}} S_0(x_t) + F \underbrace{(\eta_{ct}^{\text{QCD}}, \eta_{cc}^{\text{QCD}}, m_c, \dots)}_{\text{(NLO)}} \right)$$

BJW (90)

HN (94)

2009
2010
News

★ $\hat{B}_K \cong 0.72 \pm 0.03$

(precise and lower by
~10% vs 2007)

RBC-UKQCD
Aubin et al.
ETMC

★ $\kappa_\varepsilon \cong 0.94 \pm 0.02$
(LD Effects)

AJB + Guadagnoli (08)
+ Isidori (10)

(Nierste; Vysotsky)

Large N
 $\hat{B}_K = 0.70$

BBG (87)

★ NNLO QCD
calculation of $\eta_{cc}^{\text{QCD}}, \eta_{ct}^{\text{QCD}}$

Brod + Gorbahn (10)

(BG)

$$|\varepsilon_K^{\text{SM}}| = (1.85 \pm 0.22) \cdot 10^{-3}$$

$$|\varepsilon_{\text{exp}}| = (2.229 \pm 0.012) \cdot 10^{-3}$$

(BaBar
Belle)

using $(\sin 2\beta)_{\psi K_s} = 0.672 \pm 0.023$

(NA48, KLOE, KTeV)

Possible Solutions to ε_K - Anomaly

$$|\varepsilon_K|^{\text{SM}} \sim \kappa_\varepsilon \hat{B}_K |V_{cb}|^2 \left(\frac{1}{2} |V_{cb}|^2 R_t^2 \sin 2\beta \eta_{tt}^{\text{QCD}} S_0(x_t) + F(\eta_{ct}^{\text{QCD}}, \eta_{cc}^{\text{QCD}}, m_c, \dots) \right)$$

1.

Add New Physics to ε_K

$$\text{CMFV } S_0(x_t) \rightarrow S_0(x_t) + \Delta S_0^{\text{NP}}$$

or simply $\Delta\varepsilon_k$ (Non-MFV)

AJB
Guadagnoli

2.

$$\text{Increase } \sin 2\beta \cong 0.67 \Rightarrow 0.85$$

$$S_{\psi K_s} = \sin(2\beta + 2\varphi_{\text{NP}})$$

(Ulfit; BBGT; Ball, Fleischer;
Branco et al)

$$\varphi_{\text{NP}} \cong -8.1^\circ$$

$$\text{Large } |V_{ub}|$$

Lunghi
Soni

Super-B

3.

$$\text{Increase } R_t \rightarrow \gamma = \delta_{\text{CKM}} \approx 67^\circ \Rightarrow 82^\circ$$



LHC



4.

$$\text{Increase } |V_{cb}| \approx (41.2 \cdot 10^{-3}) \Rightarrow (43.5 \cdot 10^{-3})$$



Super-B



Diego Guadagnoli

Models investigated by TUM-Teams

(Last decade)

SM

MFV

MSSM+MFV

Z'-Models

**General
MSSM**

**Universal
Extra
Dimensions**

**RS with
custodial
protection**

**Littlest
Higgs**

**Littlest
Higgs with
T-Parity**

**SUSY+Flavour
Abelian
Symmetry
(Agashe+Carone)**

**2 Higgs
Doublet
Models**

**SUSY with
SU(3) Flavour
(Ross et al)
(RVV2)**

**SUSY with
SU(2) Flavour
(LH-currents)**

**Flavour Blind
MSSM**

4G

My Collaborators

SUSY



W. Altmannshofer

S. Gori

P. Paradisi

D. Straub

LHT



M. Blanke

B. Duling

S. Recksiegel

C. Tarantino

RS



M. Albrecht

M. Blanke

B. Duling

K. Gemmeler

S. Gori

A. Weiler

4 G



B. Duling



T. Heidsieck



C. Promberger



T. Feldmann



S. Recksiegel

2 HDM



M.V. Carlucci



S. Gori



G. Isidori

ϵ_K



D. Guadagnoli



I. Bigi



P. Ball



A. Bharucha



M. Wick



L. Calibbi




M. Nagai

4 Generations

by

4 Physics Generations



AJB



T. Feldmann



S. Recksiegel



B. Duling



C. Promberger



T. Heidsieck

5.

Models for BMFV

Most popular BSM Directions

CMFV
(constrained MFV)

MFV

(NMFV)
(GMFV)

2HDM

LHT
(Littlest Higgs
with T-parity)

SUSY
(flavour models)

Z'
(Langacker...)

RS
(Randall-Sundrum)
(Warped Extra Dimensions)

4th G

(Hou.., Soni.., Lenz.., Melic)
Munich

Vector-Like
Quarks

(Branco...,
del Aguila)



Non-Decoupling

New gauge bosons, fermions, scalars in loops
and even trees with often non-CKM interactions.

4G Model

The CKM4 matrix : New: $s_{14}, s_{24}, s_{34}, \delta_{14}, \delta_{24}, m_t, m_b, 300\text{-}600 \text{ GeV}$

$c_{12}c_{13}c_{14}$	$c_{13}c_{14}s_{12}$	$c_{14}s_{13}e^{-i\delta_{13}}$	$s_{14}e^{-i\delta_{14}}$
$-c_{23}c_{24}s_{12} - c_{12}c_{24}s_{13}s_{23}e^{i\delta_{13}}$ $-c_{12}c_{13}s_{14}s_{24}e^{i(\delta_{14}-\delta_{24})}$	$c_{12}c_{23}c_{24} - c_{24}s_{12}s_{13}s_{23}e^{i\delta_{13}}$ $-c_{13}s_{12}s_{14}s_{24}e^{i(\delta_{14}-\delta_{24})}$	$c_{13}c_{24}s_{23}$ $-s_{13}s_{14}s_{24}e^{-i(\delta_{13}+\delta_{24}-\delta_{14})}$	$c_{14}s_{24}e^{-i\delta_{24}}$
$-c_{12}c_{23}c_{34}s_{13}e^{i\delta_{13}} + c_{34}s_{12}s_{23}$ $-c_{12}c_{13}c_{24}s_{14}s_{34}e^{i\delta_{14}}$ $+c_{23}s_{12}s_{24}s_{34}e^{i\delta_{24}}$ $+c_{12}s_{13}s_{23}s_{24}s_{34}e^{i(\delta_{13}+\delta_{24})}$	$-c_{12}c_{34}s_{23} - c_{23}c_{34}s_{12}s_{13}e^{i\delta_{13}}$ $-c_{12}c_{23}s_{24}s_{34}e^{i\delta_{24}}$ $-c_{13}c_{24}s_{12}s_{14}s_{34}e^{i\delta_{14}}$ $+s_{12}s_{13}s_{23}s_{24}s_{34}e^{i(\delta_{13}+\delta_{24})}$	$c_{13}c_{23}c_{34}$ $-c_{13}s_{23}s_{24}s_{34}e^{i\delta_{24}}$ $-c_{24}s_{13}s_{14}s_{34}e^{i(\delta_{14}-\delta_{13})}$	$c_{14}c_{24}s_{34}$
$-c_{12}c_{13}c_{24}c_{34}s_{14}e^{i\delta_{14}}$ $+c_{12}c_{23}s_{13}s_{34}e^{i\delta_{13}}$ $+c_{23}c_{34}s_{12}s_{24}e^{i\delta_{24}} - s_{12}s_{23}s_{34}$ $+c_{12}c_{34}s_{13}s_{23}s_{24}e^{i(\delta_{13}+\delta_{24})}$	$-c_{12}c_{23}c_{34}s_{24}e^{i\delta_{24}} + c_{12}s_{23}s_{34}$ $-c_{13}c_{24}c_{34}s_{12}s_{14}e^{i\delta_{14}}$ $+c_{23}s_{12}s_{13}s_{34}e^{i\delta_{13}}$ $+c_{34}s_{12}s_{13}s_{23}s_{24}e^{i(\delta_{13}+\delta_{24})}$	$-c_{13}c_{23}s_{34}$ $-c_{13}c_{34}s_{23}s_{24}e^{i\delta_{24}}$ $-c_{24}c_{34}s_{13}s_{14}e^{i(\delta_{14}-\delta_{13})}$	$c_{14}c_{24}c_{34}$

Extensive New Interest in 4G

Very many papers: **Hou; Hung; Chanowitz; Novikov et al. Kribs et al.**

+

FCNC's :

Hou, Nagashima, Soddu
Soni, Alok, Giri, Mohanta, Nandi
Herrera, Benovides, Ponce
Bobrowski, Lenz, Riedl, Rohrwild
Eilam, Melic, Trampetic
AJB, Duling, Feldmann, Heidsieck, Promberger, Recksiegel
Lacker, Menzel

New Interest in Higgs-mediated FCNC's

Guidice, Lebedev (08); Agashe, Contino (09), Azatov, Toharia, Zhu (09), AJB, Gori, Duling (09); Duling (09)

Recent: Botella, Branco, Rebelo (09); Joshipura, Kodrani (08, 10)
 Pich, Tuzon (09)
 Gupta, Wells (10)

May – June
2010

- | | | |
|----------|--|---------------|
| (28 May) | Dobrescu, Fox, Martin (1005.4238) | |
| (29 May) | AJB, Carlucci, Gori, Isidori (1005.5310) | Neutral Higgs |
| (31 May) | Aranda, Montano, Ramirez-Zavaleta, Toscano, Tututi (1005.5452) | |
| (2 June) | Braeuninger, Ibarra, Simonetto | |
| (2 June) | Ligeti, Papucci, Perez, Zupan | |
| (2 June) | Jung, Pich, Tuzón | Charged Higgs |

Few Messages on Higgs-mediated FCNC's

	SUSY	2HDM	
ΔM_s	$(\tan\beta)^4$	$(\tan\beta)^2 \cdot 1/M_H^2$	
$B_{s,d} \rightarrow \mu^+ \mu^-$	$(\tan\beta)^6$	$(\tan\beta)^4 \cdot 1/M_H^4$	

Glashow
Weinberg 1977

MFV more powerful than Natural Flavour Conservation (BCGI)

General 2HDM with MFV
and flavour blind phases
(AJB, Carlucci, Gori, Isidori)



Aligned 2HDM
(Pich, Tuzón)
+ flavour blind phases

Flavour-Blind phases can be included in MFV


Mercoli, Smith (09)

Kagan, Perez, Volansky, Zupan (09) ← (could help to generate
Paradisi, Straub (09) large CP-phase in B_s -mixing)

Little Higgs Models

SUSY vs Little Higgs

Problematic quadratic divergences in m_H^2



	SUSY	Little Higgs
Quadratic divergences canceled by:	(different statistics) super-partners	(same statistics) heavy partners
Coupling relationships due to:	boson-fermion symmetry	global symmetry

The most economical in matter content: Littlest Higgs (LH)

[N. Arkani-Hamed, A.G. Cohen, E. Katz, A.E. Nelson (2002)]

valid up to $(4\pi f) \equiv \Lambda$

Original model: Arkani-Hamed, Cohen, Katz, Nelson (2002)

$$f \approx O(1\text{TeV})$$

LH

Global:

$$SU(5) \longrightarrow SO(5)$$


Local: $[SU(2) \otimes U(1)]_1 \otimes [SU(2) \otimes U(1)]_2 \longrightarrow SU(2)_L \otimes U(1)_Y$

$$(g_1) \quad (g'_1) \quad (g_2) \quad (g'_2)$$

Model with T-Parity: Cheng, Low (2003)

LHT

Theory symmetric under $[SU(2) \otimes U(1)]_1 \longleftrightarrow [SU(2) \otimes U(1)]_2$



$$g_1 = g_2 \quad g'_1 = g'_2$$

Littlest Higgs Models without and with T-Parity

New particles: (with $O(f)$ masses)

LH

Gauge Bosons: W_H^\pm, Z_H^0, A_H^0

Fermions: T

Scalars: Φ^\pm, \dots

LHT

T-even
Sector

SM Particles + T_+

T-odd
Sector

Gauge Bosons: W_H^\pm, Z_H^0, A_H^0

Fermions: T_- , Mirror Fermions

Scalars: Φ^\pm, \dots

The World of Mirror Fermions

[I. Low, hep-ph/0409025]

Required to cut-off
large 4-fermion operators
constrained by LEP

$$\begin{pmatrix} u_{1H} \\ d_{1H} \end{pmatrix} \begin{pmatrix} u_{2H} \\ d_{2H} \end{pmatrix} \begin{pmatrix} u_{3H} \\ d_{3H} \end{pmatrix}$$

Vectorial couplings under $SU(2)_L$


New Flavour Interactions
involving SM fermions,
Mirror Fermions and
 W^\pm_H, Z^0_H, A^0_H

$V_{Hu}^\dagger V_{Hd} = V_{CKM}$
[I. Low, hep-ph/0409025]
[J. Hubisz, S.J. Lee, G. Paz]


$(V_{Hu})_{ij}$ for:

Similarly for Leptons

$m_u^u = m_d^d \quad i=1,2,3$
to first order in v/f



LHT goes beyond Minimal Flavour Violation (MFV)
(without introducing new operators and non-perturbative uncertainties)
 ``visible effects in flavour physics are possible''



$$V_{Hu}^\dagger V_{Hd} = V_{CKM}$$

[Low], [Hubisz, Lee, Paz]

$$V_{Hd} = \begin{pmatrix} c_{12}^d c_{13}^d & s_{12}^d c_{13}^d e^{-i\delta_{12}^d} & s_{13}^d e^{-i\delta_{13}^d} \\ -s_{12}^d c_{23}^d e^{i\delta_{12}^d} - c_{12}^d s_{23}^d s_{13}^d e^{i(\delta_{13}^d - \delta_{23}^d)} & c_{12}^d c_{23}^d - s_{12}^d s_{23}^d s_{13}^d e^{i(\delta_{13}^d - \delta_{12}^d - \delta_{23}^d)} & s_{23}^d c_{13}^d e^{-i\delta_{23}^d} \\ s_{12}^d s_{23}^d e^{i(\delta_{12}^d + \delta_{23}^d)} - c_{12}^d c_{23}^d s_{13}^d e^{i\delta_{13}^d} & -c_{12}^d s_{23}^d e^{i\delta_{23}^d} - s_{12}^d c_{23}^d s_{13}^d e^{i(\delta_{13}^d - \delta_{12}^d)} & c_{23}^d c_{13}^d \end{pmatrix}$$

**V_{Hd} parameterization similar to CKM, but with 2 additional phases
 (the phases of SM quarks are no more free to be rotated)**
 [Blanke, AJB, Poschenrieder, Recksiegel, Tarantino, Uhlig, Weiler]

[Similar new interactions and mixing matrices appear in the lepton sector]

General Structure of the Amplitudes

LH (CMFV Model)

Non-perturbative factors

Real functions

$$A(\text{Decay}) = \sum_i B_i^{\text{SM}} \eta_{QCD}^i V_{CKM}^i F_i(m_t, m_T, m_{W_H}, \dots)$$

LHT

Real functions

$$A(\text{Decay}) = \sum_i B_i^{\text{SM}} \eta_{QCD}^i [V_{CKM}^i F_i(m_t, m_T) + V_{Hd}^i G_i(m_H^u, m_H^d, W_H^\pm, Z_H^0, A_H^0)]$$

T-even contribution: CMFV

T-odd contribution: New CP and Flavour violating Interactions
but only SM operators

LH(without T-parity) vs LHT(with T-parity)

Tree-level heavy gauge boson contributions and the triplet Φ vev make ew precision tests highly constraining
[Han, Logan, McElrath, Wang]
[Csaki, Hubisz, Kribs, Meade, Terning]

$$f \geq 2\text{-}3 \text{ TeV}$$

- The little hierarchy problem is back
- Only small effects in Flavour Physics

Buras, Poschenrieder, Uhlig, hep-ph/0410309//0501230
Buras, Poschenrieder, Uhlig, Bardeen, hep-ph/0607189
Choudhury, Gaur, Goyal, Mahajan, hep-ph/0407050
Lee, hep-ph/0408362
Fajfer, Prelovsek, hep-ph/0511048
Huo, Zhu, hep-ph/0306029
Choudhury, Gaur, Joshi, McKellar, hep-ph/0408125

These unwanted contributions are eliminated by a discrete symmetry:

T-parity

- SM particles are T-even,
- new particles are T-odd
(similarly to R-parity in SUSY)

smaller f allowed by ew tests
[Hubisz, Meade, Noble, Perelstein]

$$f \geq 500 \text{ GeV}$$

- The little hierarchy problem is solved
- Large effects are possible in Flavour Physics

General Structure of New Physics Contributions

SM : $\lambda_t^{(K)} = V_{ts}^* V_{td}$ $\lambda_t^{(d)} = V_{tb}^* V_{td}$ $\lambda_t^{(s)} = V_{tb}^* V_{ts}$

Amplitudes

: $\lambda_t^{(i)} X_{SM}(m_t)$ $\lambda_t^{(i)} Y_{SM}(m_t)$

$v\bar{v}$ in the final state $\mu^+ \mu^-$ in the final state

$i = K, B_d, B_s$

**Universality
of short
distance
functions**

LHT

: $X_i = \underbrace{X_{SM}(m_t) + \bar{X}_{even}}_{\text{real}} + \frac{1}{\lambda_t^{(i)}} \xi_i \bar{X}_{odd} \equiv |X_i| e^{i\theta_X^i}$

\uparrow
 V_{Hd}

$\underbrace{\quad}_{\text{complex}}$

$Y_i = \underbrace{Y_{SM}(m_t) + \bar{Y}_{even}}_{\text{real}} + \frac{1}{\lambda_t^{(i)}} \xi_i \bar{Y}_{odd} \equiv |Y_i| e^{i\theta_Y^i}$

(mirror fermions)

**Breakdown
of
Universality**

Natural Expectations

$$X_i = X_{SM}(m_t) + \bar{X}_{even} + \frac{1}{\lambda_t^{(i)}} \xi_i \bar{X}_{odd} \equiv |X_i| e^{i\theta_X^i}$$

(similarly for Y_i)

V_{Hd}

$i = K, B_d, B_s$

$$\frac{1}{\lambda_t^{(K)}} \approx 2 \cdot 10^3$$

$$\frac{1}{\lambda_t^{(d)}} \approx 100$$

$$\frac{1}{\lambda_t^{(s)}} \approx 25$$


$$\left\{ \begin{array}{l} \text{Natural} \\ \text{size} \\ \text{of NP} \\ \text{contributions} \end{array} \right\} : K \gg B_d > B_s$$

But can be reversed for
special structures of V_{Hd}

Randall-Sundrum Framework (Express Summary)

5D spacetime with warped metric:

$$ds^2 = e^{-2ky} \eta_{\mu\nu} dx^\mu - dy^2 \quad 0 \leq y \leq L$$



- fermions and gauge bosons live in the bulk
- Higgs localised on IR brane

(Chang, Okada et al.
Grossman, Neubert
Gherghetta, Pomarol)


- energy scales suppressed by warp factor e^{-ky}
natural solution to the gauge hierarchy problem.
- Kaluza-Klein (KK) excitations of both SM fermions and gauge bosons live close to the IR brane.

Fermion Localisation and Yukawa Couplings

SM fermion (zero mode) shape function depends strongly on bulk mass parameter characteristic for a given fermion:

$$f^{(0)}(y, c) \propto e^{\left(\frac{1}{2}-c\right)y}$$

UV brane IR brane



$c > \frac{1}{2}$: localisation near UV brane

$c < \frac{1}{2}$: localisation near IR brane

effective 4D Yukawa couplings:

$$(Y_{u,d})_{ij} = (\lambda_{u,d})_{ij} f_i^Q f_j^u$$

- $\lambda_{u,d} \sim 0(1)$ anarchic complex 3×3 matrices $\equiv Y_{5D}$
- hierarchical structure of quark masses and CKM parameters can be naturally generated by exponential suppression of $f^{Q,u,d}$ at IR brane.

Bulk Profiles of SM Gauge Bosons

- Gluons and Photon : flat (protection by Gauge symmetry)
- W^\pm, Z : flat before EWSB
but
distorted near the IR brane after EWSB $\propto \left(\frac{v^2}{M_{KK}^2}\right)$
- Equivalently : Mixing of KK gauge bosons with W^\pm, Z in the process of EWSB modifies the couplings of mass eigenstates W^\pm, Z

- Recall : All KK gauge bosons live close to the IR brane
- All KK fermions live close to the IR brane

First Implications for Phenomenology

1.

Gauge-Fermion
Interactions:
Overlaps of shape
functions



Non-universalities
in
Gauge Couplings

(in flavour)

of $\left\{ \begin{array}{l} \text{KK-gauge bosons} \\ W^\pm, Z \end{array} \right\}$
to $\{\text{SM fermions}\}$



2.

Impact on
Electroweak Precision
Observables

$SU(2)_L \otimes U(1)_Y$

S parameter : $M_{KK} \geq (2-3) \text{ TeV}$
T parameter: $M_{KK} \geq 10 \text{ TeV}$

Agashe, Delgado, May, Sundrum (2003)
Csaki, Grojean, Pilo, Terning (2003)

Also problems with $Z b_L \bar{b}_L$


3.

Tree Level FCNC mediated by KK gauge bosons and Z (breakdown of standard GIM mechanism)

$$\mathbf{d} \equiv \begin{pmatrix} \mathbf{d} \\ \mathbf{s} \\ \mathbf{b} \end{pmatrix}$$

$$\bar{\mathbf{d}} \mathbf{D}_L^+ \begin{pmatrix} \mathbf{a} & & \\ & \mathbf{b} & \\ & & \mathbf{c} \end{pmatrix} \mathbf{D}_L \gamma_\mu \mathbf{Z}^\mu \mathbf{d} \neq \bar{\mathbf{d}} \gamma_\mu \mathbf{Z}^\mu \mathbf{d}$$

(non-universality)



But RS-GIM helps in avoiding disaster.

Gherghetta, Pomarol
Huber, Shafi
Agashe, Soni, Perez

4.

Mixing of KK fermions with SM fermions and
mixing of KK gauge bosons with SM gauge bosons



Breakdown of Unitarity of the CKM matrix

5.

$\left\{ \text{Tree level exchanges of } G_{KK} \text{ and } Z \right\} \rightarrow \left\{ \text{Contributions of new operators. In particular } Q_{LR} \text{ operators in addition to } Q_{LL}, Q_{RR} \right\}$

6.

$\left\{ \text{The presence of three } 3 \times 3 \text{ hermitian bulk matrices } c^q, c^u, c^d \text{ in addition to usual Yukawa couplings} \right\} \rightarrow \left\{ \begin{array}{l} \text{New flavour and CP violating parameters:} \\ \left\{ \begin{array}{ll} 3 * 6 = 18 & \text{real} \\ 3 * 3 = 9 & \text{phases} \end{array} \right. \end{array} \right\}$

Non-MFV

A RS Model with Custodial Protection


$SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_X \otimes P_{LR}$

Gauge Group in the Bulk

$P_{LR} : SU(2)_L \leftrightarrow SU(2)_R$

P_{LR} symmetric fermion representations

A Realistic Model in the Reach of the LHC



$$SU(2)_R \times U(1)_X \rightarrow U(1)_Y$$

by boundary conditions

+ $(L \leftrightarrow R)$ -symmetric fermion representations

$$SU(2)_L \times SU(2)_R \rightarrow SU(2)_V$$

by Higgs VEV

low energy theory: $SU(2)_L \times U(1)_Y \rightarrow U(1)_{\text{em}}$

What is protected in this Model?

(up to small symmetry breaking due to UV boundry conditions)

A.

T-Parameter

Agashe, Delgado, May, Sundrum (0308036)
Csaki, Grojean, Pilo, Terning (0308038)



B.

$Z\bar{b}_L b_L$

Agashe, Contino, Rold, Pomarol (0605341)

C.

$Z\bar{d}_L^i d_L^j$

Blanke, AJB, Duling, Gori, Weiler (0809.1073)
Blanke, AJB, Duling, Gemmler, Gori (0812.3803)



D.

$Z\bar{u}_R^i u_R^j$

AJB, Duling, Gori (0905.2318)

But: $Z\bar{d}_R^i d_R^j, Z\bar{u}_L^i u_L^j, W^+ \bar{u}_L^i d_L^j$ not protected

Particle Content of the Model

Albrecht, Blanke, AJB, Duling, Gemmler (0903.2415)

Gauge sector

$$W^\pm, \quad W_H^\pm, \quad W'^\pm \\ Z^0, \quad Z_H, \quad Z'$$

KK

$$A, \quad A^{(\prime)} \\ G^a, \quad G^{a(\prime)}$$

KK

Quark sector
(i=1,2,3)

$$(2,2) = \begin{pmatrix} \chi^{u_i} (-+)_{5/3} & q^{u_i} (++)_{2/3} \\ \chi^{d_i} (-+)_{2/3} & q^{d_i} (++)_{1/3} \end{pmatrix}_L$$

$SU(2)_L \otimes SU(2)_R$

$$(3,1) = \begin{pmatrix} \psi'^i (-+)_{5/3} \\ U'^i (-+)_{2/3} \\ D'^i (-+)_{-1/3} \end{pmatrix}_R \oplus \begin{pmatrix} \psi''^i (-+)_{5/3} \\ U''^i (-+)_{2/3} \\ D^i (++)_{-1/3} \end{pmatrix}_R = (1,3)$$

+
states of
opposite
chirality

$Q=5/3$
Fermions!

(Feynman rules worked out for SM and n=1 KK modes)

6.

Patterns of Flavour Violation Beyond the SM

Three Strategies in Waiting for NP

1.

Precision Calculations

within the SM

Background to NP

$$(B \rightarrow X_s \gamma, K^+ \rightarrow \pi^+ v\bar{v}, B \rightarrow X_s l^+ l^-)$$

2.

Bottom-Top Approach

Powerful in Electroweak
Precision Physics

Personal
View

In Flavour Physics less useful due to the presence
of many operators (Buchmüller, Wyler: 1990)

Exception: Minimal Flavour Violation Hypothesis

3.

Top-Bottom Approach

Study of patterns of flavour violation in concrete
NP models. Correlations between observables !

Search for New Physics in 2010's through Flavour Physics


★ To search for NP in rare K, B_d, B_s, D decays,
CP in B_s, D decays,
Lepton Flavour Violations

★ Correlations will be crucial to distinguish various NP scenarios



Specific Plots (Correlations)

$\text{Br}(K_L \rightarrow \pi^0 \bar{v} \bar{v})$ vs $\text{Br}(K^+ \rightarrow \pi^+ \bar{v} \bar{v})$
 $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ vs $S_{\psi\phi}$ ★
 $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ vs $\text{Br}(B_d \rightarrow \mu^+ \mu^-)$
 $\text{Br}(K^+ \rightarrow \pi^+ \bar{v} \bar{v})$ vs $S_{\psi\phi}$
 d_n vs $S_{\phi K_s}$
 $A_{CP}(B \rightarrow X_S \gamma)$ vs $S_{\phi K_s}$
 $\text{Br}(\tau \rightarrow \mu \gamma)$ vs $\Delta(g-2)_\mu$
 $\text{Br}(\tau \rightarrow \mu \mu \mu)$ vs $\text{Br}(\tau \rightarrow \mu \gamma)$
 $\text{Br}(\mu \rightarrow 3e)$ vs $\text{Br}(\mu \rightarrow e \gamma)$



Patterns of Flavour Violations in specific NP Models

Superstars of 2010 – 2015 (Flavour Physics)

$$S_{\psi\phi} \\ (B_s \rightarrow \phi\phi)$$

$$B_s \rightarrow \mu^+ \mu^- \\ (B_d \rightarrow \mu^+ \mu^-) \\ (B^+ \rightarrow \tau^+ \nu_\tau)$$

$$K^+ \rightarrow \pi^+ \bar{\nu}\bar{\nu} \\ (K_L \rightarrow \pi^0 \bar{\nu}\bar{\nu}) \\ (B_d \rightarrow K^* \mu^+ \mu^-)$$

γ
from Tree
Level
Decays

$$\mu \rightarrow e\gamma \\ \tau \rightarrow \mu\gamma \\ \tau \rightarrow e\gamma \\ \mu \rightarrow 3e \\ \tau \rightarrow 3 \text{ leptons}$$

$$\varepsilon'/\varepsilon \\ (\text{Lattice}) \\ \text{EDM's} \\ (g-2)_\mu$$

Superstars enter the Scene

in the context of

SUSY

LHT

RS

4G

2HDM

(flavour models)

Number of new Flavour Parameters

(Quark Sector)

(physical)

Real

\mathcal{CP} Phases

SUSY

36

27

(R-parity)

4 G

5

2

LHT

7

3

some
sensitivity
to UV

RS

18

9

SM

9

1

Prima Donna of 2010 – Flavour Physics

Mixing Induced CP Asymmetry in $B_s \rightarrow \psi\varphi$ ($S_{\psi\varphi}$) (A_{SL}^S)

(TH very clean; ^{*)} Analog of $S_{\psi K_s}$)

$$S_{\psi\varphi} = \sin(2|\beta_s| - 2\varphi_s^{\text{new}}) \stackrel{\text{SM}}{\cong} 0.035$$

$$V_{ts} = -|V_{ts}|e^{-\beta_s} \\ (\beta_s = -1^\circ)$$

CDF : Hints for a much larger
D0 value

New Phase in $B_s^0 - \bar{B}_s^0$ mixing

Prima Donna of 2010 – Flavour Physics

Mixing Induced CP Asymmetry in $B_s \rightarrow \psi\phi$ ($S_{\psi\phi}$) (A_{SL}^s)

(TH very clean; ^{*)} Analog of $S_{\psi K_s}$)

$$S_{\psi\phi} = \sin(2|\beta_s| - 2\phi_s^{\text{new}}) \stackrel{\text{SM}}{\cong} 0.035$$

$$V_{ts} = -|V_{ts}|e^{-\beta_s} \\ (\beta_s = -1^\circ)$$

CDF : Hints for a much larger
D0 value

New Phase in $B_s^0 - \bar{B}_s^0$ mixing

Preliminary
result from
Lenz, Nierste
+ CKM fitters

(soon!)

$$S_{\psi\phi} = 0.78^{+0.12}_{-0.17}$$

3 σ : [0.07, 1]
range

Without latest
CDF result !

(0.8 σ)

Louise Oakes's talk
in Torino

But CDF cannot exclude
values above 0.5 !

Patterns of Deviations from CPV – SM Predictions

$$K^0 - \bar{K}^0 \quad (\varepsilon_K) \quad \frac{|\varepsilon_K|_{\text{SM}}}{|\varepsilon_K|_{\text{exp}}} \approx 0.83 \pm 0.10$$

$$B_d^0 - \bar{B}_d^0 \quad (S_{\psi K_s}) \quad (S_{\psi K_s}) \approx \begin{cases} 0.74 \pm 0.04 & (\text{SM}) \quad (\text{UTfit}) \\ 0.672 \pm 0.022 & (\text{exp}) \end{cases}$$

$$B_s^0 - \bar{B}_s^0 \quad (S_{\psi\phi}) \quad \frac{(S_{\psi\phi})_{\text{exp}}}{(S_{\psi\phi})_{\text{SM}}} \approx 10 - 20$$

Do these deviations signal non-MFV interactions at work ?

(non-SUSY)

General 2HDM with MFV and Flavour Blind CPV Phases

(1005.5310)

(AJB, Carlucci, Gori, Isidori)

Provides correct pattern

$$\varepsilon_K : \quad \text{Diagram} \quad \approx \left[\frac{m_d m_s}{M_H^2} \right] m_t^4 (\tan \beta)^2 (V_{ts}^* V_{td})^2 \quad (\text{tiny})$$
$$S_{\psi K_s} : \quad \text{Diagram} \quad \approx \left[\frac{m_b m_d}{M_H^2} \right] m_t^4 (\tan \beta)^2 (V_{tb}^* V_{td})^2 e^{i\phi_{\text{new}}}$$
$$S_{\psi\phi} : \quad \text{Diagram} \quad \approx \left[\frac{m_b m_s}{M_H^2} \right] m_t^4 (\tan \beta)^2 (V_{tb}^* V_{ts})^2 e^{i\phi_{\text{new}}}$$

$$S_{\psi K_s} = \sin(2\beta - \theta_d^H) \quad S_{\psi\phi} \approx \sin(\theta_s^H)$$

$$\sin 2\beta > S_{\psi K_s}$$

ZakopaneBuras2010

($|\varepsilon_K|$ enhanced)

$$\frac{\theta_d^H}{\theta_s^H} \approx \frac{m_d}{m_s} \approx \frac{1}{17}$$


$$\begin{aligned} \tan \beta &\approx 10 - 20 \\ M_H &\approx 250 \text{ GeV} \end{aligned}$$

Large
RG QCD
effects
 Q_{LR}

$S_{\psi K_s}$ vs $S_{\psi\phi}$ and $|\varepsilon_K|$ vs $S_{\psi\phi}$
in a General 2HDM with MFV and Flavour Blind CPV

(AJB, Carlucci, Gori, Isidori)

Correct pattern of NP effects





$$B_s \rightarrow \mu^+ \mu^- \text{ and } B_d \rightarrow \mu^+ \mu^-$$

Z-Penguin (SM
+ Boxes CMFV)

SM

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) = (3.2 \pm 0.2) \cdot 10^{-9}$$

$$\text{Br}(B_d \rightarrow \mu^+ \mu^-) = (1.0 \pm 0.1) \cdot 10^{-10}$$

Error dominated by $\hat{B}_{d,s}$

AJB (03)

CMFV
“Golden Relation”

$$\frac{\text{Br}(B_s \rightarrow \mu^+ \mu^-)}{\text{Br}(B_d \rightarrow \mu^+ \mu^-)} = \frac{\hat{B}_d}{\hat{B}_s} \frac{\tau(B_s)}{\tau(B_d)} \frac{\Delta M_s}{\Delta M_d}$$

$(\Delta B = 1)$ (0.95 ± 0.03) $(\Delta B = 2)$
Lattice

Valid in all CMFV models

Can be strongly violated in SUSY, LHT, RS, 4G

95% CL

$$\text{Br}(B_s \rightarrow \mu^+ \mu^-) \leq \begin{cases} 3.3 \cdot 10^{-8} & (\text{CDF}) \\ 5.3 \cdot 10^{-8} & (\text{D0}) \end{cases}$$

$$\text{Br}(B_d \rightarrow \mu^+ \mu^-) \leq 1 \cdot 10^{-8} \text{ (CDF)}$$


Zakopan

LHC should be able to discover $B_s \rightarrow \mu^+ \mu^-$ even at the SM level

Fleischer et al

$B_{s,d} \rightarrow \mu^+ \mu^-$ in Various Models

Babu, Kolda (99), ... + 100



$$Br(B_{s,d} \rightarrow \mu^+ \mu^-) \sim \frac{(\tan\beta)^6}{M_A^4}$$

Can reach CDF and DØ bounds

$$\frac{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{4G}}{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{SM}} \leq 4$$

$$\frac{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{SUSY}}{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{SM}} \leq 20$$

$$\frac{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{LHT}}{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{SM}} \leq 1.3$$

$$\frac{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{RS}}{Br(B_{s,d} \rightarrow \mu^+ \mu^-)_{SM}} \leq 1.1$$

(Z-penguin)
(Blanke et al) (09)

(Z-penguin + Z-tree with
r.h. couplings)
(Custodial protection at work)
(Gori et al) (08)


CDF, D0
LHCb

$\text{Br}(B_s \rightarrow \mu^+ \mu^-) \text{vs } S_{\psi\phi}$


SUSY

ABGPS

■ Solution 3 to ε_K -Anomaly
Abelian (AC)



■ Solution 1 to ε_K -Anomaly
Non-Abelian (RVV)




(Large Effects in $D^0-\bar{D}^0$)

(Small Effects in $D^0-\bar{D}^0$)


$$\text{Br}(\mathbf{B}_s \rightarrow \mu^+ \mu^-) \text{ vs } S_{\psi\phi}$$

4G

BDFHPR



CDF D0



Adding ε'/ε Constraint

4G has hard time to describe simultaneously ε'/ε and $S_{\psi\phi} > 0.2$ if $B_{6,8}$ within 20% from large N values



ABGPS

$\text{Br}(\mathbf{B}_d \rightarrow \mu^+ \mu^-) \text{ vs } \text{Br}(\mathbf{B}_s \rightarrow \mu^+ \mu^-)$

SUSY

MFV

AJB; Hurth, Isidori, Kamenik, Mescia





RVV2

(RH currents)

LH currents

$$B_{s,d} \rightarrow \mu^+ \mu^- \text{ in 2HDM - MFV} \approx (\tan \beta)^4 / M_A^4$$



(AJB, Carlucci, Gori, Isidori)



$$\text{Br}(B_d \rightarrow \mu^+ \mu^-) \text{ vs } \text{Br}(B_s \rightarrow \mu^+ \mu^-)$$

4G

BDFHPR



$$\text{Br}(B_d \rightarrow \mu^+ \mu^-) \text{ vs } S_{\psi\varphi}$$

Very different patterns
compared with SUSY,
2HDM, MFV

$$K^+ \rightarrow \pi^+ \nu\bar{\nu} \text{ and } K_L \rightarrow \pi^0 \nu\bar{\nu} \quad (\text{Z}^\circ\text{-penguins})$$

(TH cleanest FCNC decays in Quark Sector)

Extensive
TH efforts
over
20 years

: Buchalla, AJB; Misiak, Urban (NLO QCD)
AJB, Gorbahn, Haisch, Nierste (NNLO QCD)
Brod, Gorbahn (QED, EW two loop)
Isidori, Mescia, Smith (several LD analyses)
Buchalla, Isidori (LD in $K_L \rightarrow \pi^0 \nu\bar{\nu}$)

$$\frac{\text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu})}{\text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu})} = 3.2 \pm 0.2$$

SM

$$\text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu}) = (8.4 \pm 0.7) \cdot 10^{-11}$$

$$\text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu}) = (2.6 \pm 0.4) \cdot 10^{-11}$$

Exp

$$\text{Br}(K^+ \rightarrow \pi^+ \nu\bar{\nu}) = \left(17 \begin{array}{l} +11 \\ -10 \end{array} \right) \cdot 10^{-11}$$

$$\text{Br}(K_L \rightarrow \pi^0 \nu\bar{\nu}) \leq 6.8 \cdot 10^{-8}$$

(E787, E949 Brookhaven)

(E391a, KEK)

Future :

NA62
Project X (FNAL)

Both very
sensitive to
New Physics


CP-conserving
TH uncertainty 2-3%

J-PARC KOTO

CP-Violation in Decay
TH uncertainty 1-2%





$K_L \rightarrow \pi^0 \nu \bar{\nu}$ vs. $K^+ \rightarrow \pi^+ \nu \bar{\nu}$



$K_L \rightarrow \pi^0 \nu \bar{\nu}$ vs. $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ in 4G

BDFHPR




With ε'/ε Constraint

Much larger enhancements than
in LHT, RS, SUSYf possible

$$K^+ \rightarrow \pi^+ \nu\bar{\nu} \text{ vs. } S_{\psi\phi}$$


(Simultaneous Large Enhancements unlikely)



**Blanke, AJB, Duling,
Gori, Weiler**


**Blanke, AJB, Duling,
Recksiegel, Tarantino**

LHT



$K^+ \rightarrow \pi^+ \nu\bar{\nu}$ vs $S_{\psi\phi}$ (4G)

(Simultaneous Large Enhancements Possible)



DNA Tests of Flavour Models

O_i : *Observables*

M_i : *Models beyond SM*

	M_1	M_2	M_3	M_4	M_5
O_1	★★★	★	★	★	★★
O_2	★	★★	★★★	★★	★
O_3	★★	★★★	★★	★	★
O_4	★★★	★★	★	★★★	★★
O_5	★	★★★	★	★★	★★★



Very large New Physics effect



Moderate New Physics effect



Very small New Physics effect

DNA Tests of Flavour Models

	AC	RVV2	AKM	δLL	FBMSSM	LHT	RS	4G
$D^0 - \bar{D}^0$	★★★	★	★	★	★	★★★	?	★★
ϵ_K	★	★★★	★★★	★	★	★★	★★★	★★
$S_{\psi\phi}$	★★★	★★★	★★★	★	★	★★★	★★★	★★★
$S_{\phi K_S}$	★★★	★★	★	★★★	★★★	★	?	★★
$A_{CP}(B \rightarrow X_s \gamma)$	★	★	★	★★★	★★★	★	?	★
$A_{7,8}(B \rightarrow K^* \mu^+ \mu^-)$	★	★	★	★★★	★★★	★★	?	★★
$A_9(B \rightarrow K^* \mu^+ \mu^-)$	★	★	★	★	★	★	?	★★
$B \rightarrow K^{(*)} \nu \bar{\nu}$	★	★	★	★	★	★	★	★
$B_s \rightarrow \mu^+ \mu^-$	★★★	★★★	★★★	★★★	★★★	★	★	★★★
$K^+ \rightarrow \pi^+ \nu \bar{\nu}$	★	★	★	★	★	★★★	★★★	★★★
$K_L \rightarrow \pi^0 \nu \bar{\nu}$	★	★	★	★	★	★★★	★★★	★★★
$\mu \rightarrow e \gamma$	★★★	★★★	★★★	★★★	★★★	★★★	★★★	★★★
$\tau \rightarrow \mu \gamma$	★★★	★★★	★	★★★	★★★	★★★	★★★	★★★
$\mu + N \rightarrow e + N$	★★★	★★★	★★★	★★★	★★★	★★★	★★★	★★★
d_n	★★★	★★★	★★★	★★	★★★	★	★★★	★
d_e	★★★	★★★	★★	★	★★★	★	★★★	★
$(g-2)_\mu$	★★★	★★★	★★	★★★	★★★	★	?	★


2020 Vision

	NEW SM
$D^0 - \bar{D}^0$	★★
ϵ_K	★★
$S_{\psi\phi}$	★★★
$S_{\phi K_S}$	★★
$A_{\text{CP}}(B \rightarrow X_s \gamma)$	★
$A_{7,8}(B \rightarrow K^* \mu^+ \mu^-)$	★★
$A_9(B \rightarrow K^* \mu^+ \mu^-)$	★
$B \rightarrow K^{(*)} \nu \bar{\nu}$	★★★
$B_s \rightarrow \mu^+ \mu^-$	★★★
$K^+ \rightarrow \pi^+ \nu \bar{\nu}$	★★
$K_L \rightarrow \pi^0 \nu \bar{\nu}$	★★★
$\mu \rightarrow e \gamma$	★★★
$\tau \rightarrow \mu \gamma$	★★★
$\mu + N \rightarrow e + N$	★★★
d_n	★★★
d_e	★★★
$(g-2)_\mu$	★★

7.

Outlook

In our search for a more
fundamental theory we need
to improve our understanding
of Flavour



Flavour Physics (Quarks and Leptons)

Final Messages of this Talk


Many observables (decays) not measured yet or measured poorly. Flavour Physics only now enters the precision era.

:



Spectacular deviations from SM still possible

Interplay



Direct searches at Tevatron, LHC, ILC

Flavour Physics (Quarks and Leptons)

DNA Flavour Test of NP models

Final Messages of this Talk

:

Many observables (decays) not measured yet or measured poorly. Flavour Physics only now enters the precision era.



Spectacular deviations from SM still possible

Interplay

Direct searches at Tevatron, LHC, ILC

Correlations between various observables can distinguish NP scenarios easier than LHC !



Great discoveries and goals are just ahead of us !

Superstars of 2010 – 2015 (Flavour Physics)

$S_{\psi\phi}$
 $(B_s \rightarrow \phi\phi)$



$B_s \rightarrow \mu^+ \mu^-$
 $(B_d \rightarrow \mu^+ \mu^-)$

$K^+ \rightarrow \pi^+ \bar{v}v$
 $(K_L \rightarrow \pi^0 \bar{v}v)$

$(B^+ \rightarrow \tau^+ \bar{v}_\tau)$


$(B_d \rightarrow K^* \mu^+ \mu^-)$

γ
from Tree
Level
Decays

$\mu \rightarrow e\gamma$
 $\tau \rightarrow \mu\gamma$
 $\tau \rightarrow e\gamma$
 $\mu \rightarrow 3e$
 $\tau \rightarrow 3 \text{ leptons}$

ϵ'/ϵ
(Lattice)

EDM's
 $(g-2)_\mu$



Construction of
New Models

Phenomenology
of Lepton Flavour
Physics

Theory of
Quark Masses
and Mixing Angles

Electric Dipole
Moments and
 $(g-2)\mu$

Flavour Violation
in High Energy
Collisions

Dark Matter and
Baryogenesis

Grand Unification
Theories

New Techniques
for Electroweak
Symmetry Breaking

Theory of
Lepton Masses and
Mixing Angles

New
Directions


New
Concepts

12
11
10
9
8
7
6
5
4
3
2
1

Thank
You!

Backup

ε'/ε in the Standard Model



$$\frac{\varepsilon'}{\varepsilon} = 10^{-4} \left[\frac{\text{Im } \lambda_t}{1.20 \cdot 10^{-4}} \right] F(m_t, \Lambda_{\overline{\text{MS}}}^{(4)}, m_s, B_6, B_8, \Omega_{\text{IB}})$$

$$F \approx 16 \cdot \left[\frac{110 \text{ MeV}}{m_s(2 \text{ GeV})} \right]^2 \left[B_6 (1 - \Omega_{\text{IB}}) - \tilde{Z}(m_t) B_8 \right] \left(\frac{\Lambda_{\overline{\text{MS}}}^{(4)}}{340 \text{ MeV}} \right)$$

$$\tilde{Z}(m_t) \cong 0.4 \left[\frac{m_t}{165 \text{ GeV}} \right]^{2.5}; \quad \Omega_{\text{IB}} = \text{Isospin Breaking}$$

$$\text{Im } \lambda_t = \text{Im} (V_{ts}^* V_{td}) = |V_{ub}| |V_{cb}| \sin \delta$$

Basic
Parameters

: $\text{Im } \lambda_t, \Lambda_{\overline{\text{MS}}}^{(4)}, B_6, B_8, m_s, \Omega_{\text{IB}}$

First Round of Measurements

$$\frac{\varepsilon'}{\varepsilon} = \begin{cases} (23 \pm 6.5) \cdot 10^{-4} & (\text{NA31}) \\ (7.4 \pm 5.9) \cdot 10^{-4} & (\text{E731}) \end{cases}$$

Second Round of Measurements

$$\frac{\varepsilon'}{\varepsilon} = \begin{cases} (14.7 \pm 2.2) \cdot 10^{-4} & (\text{NA48}) \\ (20.7 \pm 2.8) \cdot 10^{-4} & (\text{KTeV}) \end{cases}$$

Grand
Average

:

$$\frac{\varepsilon'}{\varepsilon} = (16.6 \pm 1.6) \cdot 10^{-4}$$

Waiting for KLOE

Direct CP Violation
firmly established



Starting Point

:

$$\mathcal{L} = \mathcal{L}_{\text{SM}}(g_i, m_i, V_{\text{CKM}}^i) + \mathcal{L}_{\text{NP}}(g_i^{\text{NP}}, m_i^{\text{NP}}, V_{\text{NP}}^i)$$

Goal


:

Identify the effects of \mathcal{L}_{NP} in weak decays
in the presence of the background from \mathcal{L}_{SM}


First Implication from \mathcal{L}

:


Feynman Diagrams



$B_d^0 - \bar{B}_d^0$ Mixing



$K^+ \rightarrow \pi^+ \nu \bar{\nu}, \quad K_L \rightarrow \pi^0 \nu \bar{\nu}$



+ NP

Putting $SO(10)$ -SUSY-GUT of Dermisek-Raby into difficulties

M. Albrecht, W. Altmannshofer, AJB, D. Guadagnoli, D. Straub

1.

The Model gives a nice description of quark and lepton masses, PMNS and most of CKM elements.

Also
SUSY
Spectrum

2.

But fails to describe simultaneously the data on

$$B_{s,d} \rightarrow \mu^+ \mu^-, B \rightarrow X_s \gamma, B \rightarrow X_s l^+ l^-, B_u \rightarrow \tau \nu$$

3.

Gives $|V_{ub}| \approx 3.2 \cdot 10^{-3}$

$$< \underbrace{(4.2 \pm 0.3) \cdot 10^{-3}}_{\text{Exp.}}$$



Generally
too low

Some recent
solutions:
Altmannshofer et al.

Very strong Constraints on New Physics

$$\text{Br}(B \rightarrow X_s \gamma)_{\text{exp}} = (3.52 \pm 0.24) \cdot 10^{-4}$$

$$\text{Br}(B \rightarrow X_s \gamma)_{\text{SM}} = \begin{cases} (3.15 \pm 0.23) \cdot 10^{-4} & (\text{Misiak et al}) \\ (2.98 \pm 0.26) \cdot 10^{-4} & (\text{Becher, Neubert}) \end{cases}$$

$$\text{Br}(B \rightarrow X_s l^+ l^-)_{\text{exp}} = \begin{cases} (1.6 \pm 0.5) \cdot 10^{-6} & (\text{low } q^2) \\ (4.4 \pm 1.3) \cdot 10^{-7} & (\text{high } q^2) \end{cases}$$

$$\text{Br}(B \rightarrow X_s l^+ l^-)_{\text{SM}} = \begin{cases} (1.6 \pm 0.1) \cdot 10^{-6} & (\text{low } q^2) \\ (2.3 \pm 0.8) \cdot 10^{-6} & (\text{high } q^2) \end{cases}$$

$$A_{CP}(B \rightarrow X_s \gamma)_{\text{exp}} = 0.004 \pm 0.036$$

$$A_{CP}(B \rightarrow X_s \gamma)_{\text{SM}} = 0.004 \pm 0.002$$

(Still factor 10 enhancement possible !)

Isidori et al. (incl.)
Gorbahn et al. (incl.)
Feldmann et al. (excl.)

Zero in A_{FB}

$$\hat{s}_0 = (3.50 \pm 0.12) \text{GeV}^2$$



TH
very clean

All this can be improved
at Super-B
Super-Belle

$$B^+ \rightarrow \tau^+ \nu$$

$$\text{Br}(B^+ \rightarrow \tau^+ \nu)_{\text{exp}} = (1.4 \pm 0.4) \cdot 10^{-4} \quad (\text{Belle, BaBar})$$

$$\text{Br}(B^+ \rightarrow \tau \nu)_{\text{SM}} \approx G_F^2 F_B^2 |V_{ub}|^2 = (0.95 \pm 0.20) \cdot 10^{-4}$$

$$\frac{\text{Br}(B^+ \rightarrow \tau \nu)_{\text{MSSM}}}{\text{Br}(B^+ \rightarrow \tau \nu)_{\text{SM}}} = \left[1 - \left(\frac{m_B}{m_{H^\pm}} \right)^2 \frac{\tan^2 \beta}{1 + \varepsilon_0 \tan \beta} \right]^2$$

(Hou)
(Isidori, Paradisi)

This decay could be problematic for
MSSM-MFV with large $\tan \beta$

Tree-Level
 H^+ exchange

Altmannshofer, AJB, Guadagnoli, Wick (07)

The General Mechanism of Little Higgs Models

The “little Higgs” is a pseudo-Nambu-Goldstone boson of a spontaneously broken symmetry. This symmetry is also explicitly broken but only “collectively”, i.e. the symmetry is broken when two or more couplings in the Lagrangian are non-vanishing. Setting any one of these couplings to zero restores the symmetry and therefore the masslessness of the “little Higgs”.

[N. Arkani-Hamed, A.G. Cohen, H. Georgi (2001)]

1. The **light Higgs** is interpreted as a **Goldstone boson** of a spontaneously broken global symmetry (**G**)
2. **Gauge and Yukawa couplings** of the Higgs are introduced by **gauging a subgroup of G**
3. ``Dangerous'' quadratic corrections are **avoided at one-loop** through **Collective Symmetry Breaking**
(the Higgs becomes massive only when two couplings are non-vanishing)

- The Higgs dynamics is described (similarly to ChPT) by a **non-linear sigma model up to $\Lambda \sim 10\text{TeV}$**
- The **UV completion is unknown** (another LH?, SUSY?, ED?)

Maximal Enhancements of $S_{\psi\phi}$, $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$ and $K^+ \rightarrow \pi^+ \bar{\nu}\nu$

(without taking correlation between them)

Model	Upper Bound on ($S_{\psi\phi}$)	Enhancement of $\text{Br}(B_s \rightarrow \mu^+ \mu^-)$	Enhancement of $\text{Br}(K^+ \rightarrow \pi^+ \bar{\nu}\nu)$
CMFV	0.04	20%	20%
MFV	0.04	1000%	30%
LHT	0.30	30%	150%
RS	0.75	10%	60%
4G	0.80	400%	300%
AC	0.75	1000%	2%
RVV	0.50	1000%	10%

Large
RH Currents

RS = RS with custodial protections

AC = Agashe, Carone

RVV = Ross, Velaso-Sevilla, Vives (04)

$U(1)_F$
 $SU(3)_F$

Lepton Flavour Violation, $\Delta(g - 2)_\mu$ and EDM's

$$S_{\phi K_s} = 0.44 \pm 0.17 \quad (S_{\phi K_s})_{\text{SM}} \approx (S_{\psi K_s})_{\text{SM}} + 0.02 \approx 0.70$$

(Beneke)

(MEGA) $\text{Br}(\mu \rightarrow e\gamma) < 1.2 \cdot 10^{-11}$ \rightarrow 10^{-13} (MEG) SM: 10^{-54}

$$(a_\mu)_{\text{SM}} < (a_\mu)_{\text{exp}} \quad (3.1\sigma)$$


$$a_\mu = \frac{1}{2} (\Delta g - 2)_\mu$$

(Regan et al) $d_e < 1.6 \cdot 10^{-27}$ \rightarrow 10^{-31} $(d_e)_{\text{SM}} \approx 10^{-38}$

[e cm]

(Baker et al) $d_n < 2.9 \cdot 10^{-26}$ \rightarrow 10^{-28} $(d_n)_{\text{SM}} \approx 10^{-32}$

Simultaneous Solution to $S_{\phi K_S}$ and $S_{\psi\phi}$ Anomalies in 4G Model



$\mu \rightarrow e\gamma$: State of the Art

- ◆ **SM (+ Dirac v_R):**

very much suppressed due to the smallness of m_ν

$$Br(\mu \rightarrow e\gamma)_{SM} \approx 10^{-54}$$

- ◆ **Experimental bound:**

[MEGA Collaboration]

$$Br(\mu \rightarrow e\gamma)_{exp} < 1.2 \cdot 10^{-11} \quad (90\% C.L.)$$

It will be improved to $\sim 10^{-13}$ by MEG in 2008

- ◆ **MSSM and LHT could explain such high values.**
WED too (Agashe et al.)

Other interesting Processes

- ◆ $\mu^- \rightarrow e^- e^+ e^-$: even more constrained than $\mu \rightarrow e\gamma$

$$Br(\mu^- \rightarrow e^- e^+ e^-)_{exp} < 1.0 \cdot 10^{-12}$$

[SINDRUM Collaboration]

- ◆ $\tau \rightarrow \mu\gamma$ and $\tau \rightarrow e\gamma$: similar to $\mu \rightarrow e\gamma$

$$Br(\tau \rightarrow \mu\gamma)_{exp} < 1.6 \cdot 10^{-8}$$

[Belle, BaBar]

$$Br(\tau \rightarrow e\gamma)_{exp} < 9.4 \cdot 10^{-8}$$

[BaBar, Belle]

- ◆ $\tau \rightarrow \mu\pi$: semileptonic decay

$$Br(\tau \rightarrow \mu\pi)_{exp} < 5.8 \cdot 10^{-8}$$

(Future:
Super B)

[Belle, BaBar]

- ◆ $\mu \rightarrow e$ conversion

$$R(\mu T_i \rightarrow e T_i) < 4.3 \cdot 10^{-12}$$

10⁻¹⁸ (J-Parc)


- ◆ $K_L \rightarrow \mu e$: flavour violating in both quark and lepton sectors

$$Br(K_L \rightarrow \mu e)_{exp} < 4.7 \cdot 10^{-12}$$

[BNL E871 Collaboration]


Simultaneous Solution to Δa_μ and $S_{\psi\phi}$ Anomalies

■ **Solution 3 to ε_K -Anomaly
Abelian (AC)**





(Large Effects in $D^0-\bar{D}^0$)

■ **Solution 1 to ε_K -Anomaly
Non-Abelian (RVV)**



(Small Effects in $D^0-\bar{D}^0$)

Correlations in the SU(3) Flavour Model (RVV2)



Clear Distinction between MSSM and LHT

MSSM

$$\frac{\text{Br}(\mu^- \rightarrow e^- e^+ e^-)}{\text{Br}(\mu^- \rightarrow e^- \gamma)} \approx \frac{1}{161}$$

LHT

$$0.02 - 1$$

Both
can
reach
MEGA's
 $\mu \rightarrow e\gamma$
bound

$$\frac{\text{Br}(\tau^- \rightarrow \mu^- \mu^+ \mu^-)}{\text{Br}(\tau^- \rightarrow \mu^- \gamma)} \approx \frac{1}{435}$$

$$0.04 - 0.4$$

MSSM

: (Ellis, Hisano, Raidal, Shimizu; Arganda, Herrero; Paradisi)
(Brignole, Rossi)

LHT

: (Blanke, AJB, Duling, Poschenrieder, Tarantino) (2007)
del Aguila, Illana, Jenkins (2008), Goto, Okada, Yamamoto (2009)