

CP violation in the extended Standard Model with a complex singlet

NEDA DARVISHI

University of Warsaw

Faculty of Physics

In collaboration with MARIA KRAWCZYK

Content

- Motivation
- Lagrangian
- U(1) symmetry
- Mass matrix
- CP violation of the model
- Conclusion

Motivation

- The CP violating of SM insufficient to explain the baryon asymmetry in the Universe
- Additional sources of CP-violation is needed. We will consider a extension of the SM with a complex singlet.
- Extension of the SM with a complex singlet and complex VEV \rightarrow additional CP violating phase (SM+CS)
- IDM
- SM+CS is a part of the IDMS (Inert Doublet Model plus Complex Singlet) Bonilla, Sokolowska, Diaz-Cruz, Krawczyk and ND, arXiv:1412.8730.
 - ▶ We consider a scenario according to which the SM-like Higgs particle, comes mostly from the SM-like doublet, with a small modification coming from the singlet.
 - ▶ The inert doublet is responsible for a dark matter in agreement with data
 - ▶ There is possibility for strong enough first order phase transition at the same time spontaneous CP violation and it is important for baryogenesis.

The Model

- The model contain SM-like doublet Φ and a complex singlet χ with complex VEV
- Lagrangian

$$\mathcal{L} = \mathcal{L}_{gf}^{SM} + \mathcal{L}_Y(\psi_f, \Phi) + \mathcal{L}_{scalar}, \quad \mathcal{L}_{scalar} = T - V$$

- ▶ $\mathcal{L}_{gf}^{SM} \rightarrow$ gauge boson-fermion interaction as in the SM.
 - ▶ $\mathcal{L}_Y(\psi_f, \Phi) \rightarrow$ only Φ couple to SM fermions.
- The scalar fields

$$\Phi = \begin{pmatrix} \phi^+ \\ \frac{1}{\sqrt{2}}(v + \phi_1 + i\phi_4) \end{pmatrix}, \quad \chi = \frac{1}{\sqrt{2}}(ae^{i\xi} + \phi_2 + i\phi_3).$$

$$\langle \Phi \rangle = \frac{1}{\sqrt{2}}v \quad \text{and} \quad \langle \chi \rangle = \frac{1}{\sqrt{2}}ae^{i\xi}$$

- Kinetic term of scalar sector

$$T = (D_\mu \Phi)^\dagger (D^\mu \Phi) + \partial_\mu \chi^* \partial^\mu \chi,$$

D_μ is the covariant derivative for $SU(2) \times U(1)_Y$ gauge group

$$D_\mu = \partial_\mu - igW_\mu^a t^a - ig' Y_\phi B_\mu$$

Potential

$$V = V_{\Phi} + V_{\chi} + V_{\Phi\chi}$$

- SM term

$$V_{SM} = -\frac{1}{2}m_{11}^2\Phi^\dagger\Phi + \frac{1}{2}\lambda(\Phi^\dagger\Phi)^2.$$

- Singlet term

$$\begin{aligned}V_S = & -\frac{m_s^2}{2}\chi^*\chi - \frac{m_4^2}{2}(\chi^{*2} + \chi^2) \\ & + \lambda_{s1}(\chi^*\chi)^2 + \lambda_{s2}(\chi^*\chi)(\chi^{*2} + \chi^2) + \lambda_{s3}(\chi^4 + \chi^{*4}) \\ & + \kappa_1(\chi + \chi^*) + \kappa_2(\chi^3 + \chi^{*3}) + \kappa_3(\chi(\chi^*\chi) + \chi^*(\chi^*\chi)).\end{aligned}$$

- Singlet and Doublet interaction

$$\begin{aligned}V_{SS} = & \Lambda_1(\Phi^\dagger\Phi)(\chi^*\chi) + \Lambda_2(\Phi^\dagger\Phi)(\chi^{*2} + \chi^2) \\ & + \kappa_4(\Phi^\dagger\Phi)(\chi + \chi^*).\end{aligned}$$

Constrained Model

To simplify the model we use $U(1)$ symmetry

$$U(1) : \Phi \rightarrow \Phi, \chi \rightarrow e^{i\alpha}\chi.$$

$\langle \chi \rangle$ spontaneous breaking symmetry \rightarrow To avoid massless Nambu-Goldstone scalar $U(1)$ -soft-breaking terms

- ① $U(1)$ -symmetric terms: $m_{11}^2, m_s^2, \lambda, \lambda_{s1}, \Lambda_1,$
- ② $U(1)$ -soft-breaking terms: $m_4^2, \kappa_{2,3},$
- ③ $U(1)$ -hard-breaking terms: $\lambda_{s2}, \lambda_{s3}, \Lambda_2.$

$$V = -\frac{1}{2}m_{11}^2\Phi^\dagger\Phi + \frac{1}{2}\lambda(\Phi^\dagger\Phi)^2 - \frac{m_s^2}{2}\chi^*\chi + \lambda_{s1}(\chi^*\chi)^2 + \Lambda_1(\Phi^\dagger\Phi)(\chi^*\chi) \\ - \frac{m_4^2}{2}(\chi^{*2} + \chi^2) + \kappa_2(\chi^3 + \chi^{*3}) + \kappa_3[\chi(\chi^*\chi) + \chi^*(\chi^*\chi)].$$

Constrained Potential

$$V = -\frac{1}{2}m_{11}^2\Phi^\dagger\Phi + \frac{1}{2}\lambda(\Phi^\dagger\Phi)^2 - \frac{m_s^2}{2}\chi^*\chi + \lambda_{s1}(\chi^*\chi)^2 + \Lambda_1(\Phi^\dagger\Phi)(\chi^*\chi) - \frac{m_4^2}{2}(\chi^{*2} + \chi^2) + \kappa_2(\chi^3 + \chi^{*3}) + \kappa_3[\chi(\chi^*\chi) + \chi^*(\chi^*\chi)].$$

- Parameters $\rightarrow m_{11}^2, m_s^2, m_4^2, \lambda_{s1}, \lambda, \Lambda_1, \kappa_2, \kappa_3$.

If real \rightarrow **No explicit CP violation**

- Vacuum with spontaneous CP violation

$$\langle\Phi\rangle = \frac{1}{\sqrt{2}}v \quad \text{and} \quad \langle\chi\rangle = \frac{1}{\sqrt{2}}ae^{i\xi}$$

- **Spontaneous CP violation** \rightarrow relevant parameters $m_4^2, \kappa_2, \kappa_3$
- Positivity conditions

$$\lambda > 0,$$

$$\lambda_{s1} > 0,$$

$$\bar{\lambda}_{1S} = \Lambda_1 + \sqrt{2\lambda\lambda_{s1}} > 0.$$

Mass matrix

The mass matrix that describes the singlet-doublet mixing, in the basis of neutral fields ϕ_1, ϕ_2, ϕ_3 :

$$M = \begin{pmatrix} M_{11} & M_{12} & M_{13} \\ M_{21} & M_{22} & M_{23} \\ M_{31} & M_{32} & M_{33} \end{pmatrix}$$

$$M_{11} = v^2 \lambda$$

$$M_{12} = va\Lambda_1 \cos \xi$$

$$M_{13} = va\Lambda_1 \sin \xi$$

$$M_{22} = \frac{a^2}{2 \cos \xi} (3\sqrt{2}\kappa_2 + \sqrt{2}\kappa_3(1 + 2 \cos 2\xi) + 4\lambda_{s_1} \cos^2 \xi)$$

$$M_{23} = a^2(-3\sqrt{2}\kappa_2 + \sqrt{2}\kappa_3 + 2\lambda_{s_1} \cos \xi) \sin \xi$$

$$M_{33} = 2a^2\lambda_{s_1} \sin^2 \xi$$

Mass eigenstate

$$\begin{pmatrix} h_1 \\ h_2 \\ h_3 \end{pmatrix} = R \begin{pmatrix} \phi_1 \\ \phi_2 \\ \phi_3 \end{pmatrix}.$$

$R \rightarrow$ The rotation matrix $R = R_1 R_2 R_3$ ($c_i = \cos \alpha_i, s_i = \sin \alpha_i$):

$$R_1 = \begin{pmatrix} c_1 & s_1 & 0 \\ -s_1 & c_1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad R_2 = \begin{pmatrix} c_2 & 0 & s_2 \\ 0 & 1 & 0 \\ -s_2 & 0 & c_2 \end{pmatrix}, \quad R_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_3 & s_3 \\ 0 & -s_3 & c_3 \end{pmatrix}.$$

$$R = R_1 R_2 R_3 = \begin{pmatrix} R_{11} & R_{12} & R_{13} \\ R_{21} & R_{22} & R_{23} \\ R_{31} & R_{32} & R_{33} \end{pmatrix}.$$

The composition of h_1 in ϕ_1, ϕ_2, ϕ_3

$$h_1 = c_1 c_2 \phi_1 + (c_3 s_1 - c_1 s_2 s_3) \phi_2 + (c_1 c_3 s_2 + s_1 s_3)$$

$$\phi_1 = c_1 c_2 h_1 - c_2 s_1 h_2 - s_2 h_3.$$

Physical mass

Diagonalization of M_{mix}^2 gives the physical mass

$$RM_{mix}^2R^T = \text{diag}(M_{h_1}^2, M_{h_2}^2, M_{h_3}^2)$$

Physical higgs masses

$$M_1^2 \simeq v^2 \lambda \quad \rightarrow \text{The SM-like Higgs boson mass, 125 GeV}$$

$$M_{2,3}^2 = \frac{1}{2}(M_{22} + M_{33} \mp \sqrt{(M_{22} + M_{33})^2 + 4M_{23}^2})$$

Mass matrix for real VEV

- $\xi = 0 \rightarrow$ real VEV \rightarrow CP conserving

$$M = \begin{pmatrix} M_{11} & M_{12} & 0 \\ M_{21} & M_{22} & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$M_{11} = v^2 \lambda$$

$$M_{12} = va\Lambda_1$$

CP even state

$$M_{22} = \frac{a^2}{2}(3\sqrt{2}\kappa_2 + 3\sqrt{2}\kappa_3 + 4\lambda_{s_1})$$

Extremum conditions

- The complex singlet VEV can be rewritten as

$$\langle \chi \rangle = \frac{1}{\sqrt{2}} a e^{i\xi} = \underbrace{\frac{1}{\sqrt{2}} a \cos \xi}_{a_1} + i \underbrace{\frac{1}{\sqrt{2}} a \sin \xi}_{a_2}; \quad a^2 = a_1^2 + a_2^2$$

$$\left. \frac{\partial V}{\partial \Phi} \right|_{\substack{\langle \phi_1 \rangle \\ \langle \chi \rangle \\ \langle \chi^* \rangle}} = 0,$$

$$\left. \frac{\partial V}{\partial \chi} \right|_{\substack{\langle \phi_1 \rangle \\ \langle \chi \rangle \\ \langle \chi^* \rangle}} = 0,$$

$$\left. \frac{\partial V}{\partial \chi^*} \right|_{\substack{\langle \phi_1 \rangle \\ \langle \chi \rangle \\ \langle \chi^* \rangle}} = 0$$

Extremum conditions

Set of three equation:

$$\textcircled{1} \quad -2m_{11}^2 + 2\lambda v^2 + \Lambda_1 a^2 = 0$$

$$\textcircled{2} \quad a_1(-m_s^2 - 2m_4^2 + v^2\Lambda_1 + 2a^2\lambda_{s1}) + 3\sqrt{2}(a_1^2 - a_2^2)a\kappa_2 + a\sqrt{2}(3a_1^2 + a_2^2)\kappa_3 = 0$$

$$\textcircled{3} \quad a_2(-m_s^2 + 2m_4^2 + v^2\Lambda_1 + 2a^2\lambda_{s1} - 2\sqrt{2}aa_1(3\kappa_2 + \kappa_3)) = 0.$$

From (2) and (3) and $a_1, a_2 \neq 0$

$$\Rightarrow \quad -4m_4^2 a_1 a + 3R_2(3a_1^2 - a_2^2) + R_3 a^2 = 0 \quad \rightarrow \text{CP violation}$$

The parameters (R_2, R_3) with dimension $[M]^2$ are:

$$R_2 = \sqrt{2}a\kappa_2$$

$$R_3 = \sqrt{2}a\kappa_3$$

Region for possible CP violation


- m_4^2, R_2, R_3, ξ ($a_1^2 \geq 0, a_2^2 > 0$)

$$-4m_4^2 a_1 a_2 + 3R_2(3a_1^2 - a_2^2) + R_3 a_2^2 = 0$$

For better understanding

$$-4m_4^2 \cos \xi + 3R_2(1 + 2 \cos 2\xi) + R_3 = 0$$


Figure: (R_2, R_3, ξ) , CP violation region for fix value of m_4^2



Region for CP violation

$$-4m_4^2 \cos \xi + 3R_2(1 + 2 \cos 2\xi) + R_3 = 0$$

Figure: (R_2, R_3) , CP violation is possible in green part




$$R_2 = 0$$

$$-4m_4^2 + R_3 \cos \xi = 0$$

$$-1 < \cos \xi < 1 \rightarrow -1 < \frac{R_3}{4m_4^2} < 1$$


Figure: (m_4^2, R_3) CP violation is possible in green region



$$R_3 = 0$$

$$-4m_4^2 \cos \xi + 3R_2(1 + 2 \cos 2\xi) = 0$$

Figure: (m_4^2, R_2) , CP violation is possible in green region



Conclusion

- This model contains a $SU(2)$ doublet as in the SM and a complex singlet with a complex VEV.
- This model provide source of spontaneous CP violation
- At least one cubic term is needed in order to have CP violation in the model.
- The analysis of this simple model was performed as a part of full analyzes of the IDMS model which was confronted with LHC data for 125 GeV, precision data STU as well as astro data on dark matter .

