



Probing hadron-quark mixed phase in twin stars using f-mode


Bikram Keshari Pradhan

In collaboration with

Prof. Debarati Chatterjee (Inter University Centre for Astronomy and Astrophysics, Pune, India)

Prof. David Edwin A. Castillo (The Henryk Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences, Kraków, Poland)

[arXiv:2309.08775](https://arxiv.org/abs/2309.08775)




63. Cracow School of Theoretical Physics
Nuclear Matter at Extreme Densities and High Temperatures
September 17-23, 2023, Zakopane, Tatra Mountains, Poland




Introduction: Neutron Stars

- Stellar remnant : after the death of a 8 to 20 M_{\odot} star.
- Core CS density $\sim 5\text{-}10 \times$ nuclear density .
- Observed with electromagnetic, GW detectors.



Credit: [NASA](#)




The figure illustrates the thin atmosphere, the outer and inner crust, and the outer and inner core, with the respective densities at different depths. Adapted with permission from NASA, NICER Team.

Pc: [Nature Reviews Physics](#) volume 4, pages 237–246 (2022)

Introduction: Neutron Stars


Theoretical Models :

- Microscopic Models : based on nucleon-nucleon interactions.
- Phenomenological : model parameters are fitted to available data from nuclear and hyper-nuclear experiments.




The figure illustrates the thin atmosphere, the outer and inner crust, and the outer and inner core, with the respective densities at different depths. Adapted with permission from NASA, NICER Team.

Pc: [Nature Reviews Physics](#) volume 4, (2022)




Credit: Menezes, [Universe](#) 2021, 7(8), 267

The EoS : $P = P(\epsilon)$




TOV Equations:



$$\frac{dm}{dr} = 4\pi r^2 \epsilon(r)$$








$$\frac{dP}{dr} = - [P(r) + \epsilon(r)] \frac{m(r) + 4\pi r^3 P(r)}{r(r - 2m(r))}$$

NS Configuration:



[Pradhan et al, Phys. Rev. D 107, 023010 \(2023\).](#)





Introduction: Neutron star and Observational constraints





- GW events: GW170817, GW190425
- Post-Merger yet to be detected.

Credit: Zack Carson, Thesis


Introduction: Neutron star and Observational constraints




PC: NASA



P. Landry et al, [Phys. Rev. D. 101, 123007 \(2020\)](#).




Credit: Zack Carson, Thesis




PC: B. P. Abbott et al., [Phys. Rev. Lett. 121, 161101 \(2018\)](#), LVC

- GW events: GW170817, GW190425
- Post-Merger yet to be detected.

Neutron Star and Gravitational Wave




Credit: C. Hanna and B. Owen



Credit: CERN/Indico

- Non-radial fluid QNMs.
 - fundamental (f) mode,
 - no node, probe for mean density, (1 kHz < f < 3kHz)
 - pressure (p) mode,
 - Sound speed, (5 kHz < f < 10kHz)
 - gravity (g) mode,
 - (50 Hz < f < 500 Hz)
- R-mode, for rotating stars only.
 - Viscosity, (0.5 kHz < f < 2kHz)
- Space-time (w) mode.
 - 5 kHz < f






PC: cosmicexplorer.org/sensitivity

CS EoS and Twin Stars:

- ❖ Hadron-Quark phase transition inside the NS core
 - The puzzle: Strong or Crossover ?

- ❖ Maxwell construction with Seidov condition (Z. Seidov, 1971, [Soviet Ast., 15, 347](#)),

$$\frac{\Delta \varepsilon}{\varepsilon_c} \geq \frac{1}{2} + \frac{3 P_c}{2 \varepsilon_c}$$



[N. K. Glendenning, PRD, 4, 46, \(1992\)](#)

The EoS Model : Phenomenological Description

- [A. Ayriyan, H. Grigorian, EPJ Web of Conferences, p. 03003, \(2018\).](#)
- [A. Ayriyan, N. Bastian, D. Blaschke, H. Grigorian, K. Maslov, D. N. Voskresensky, PRC 97, 045802 \(2018\).](#)
- [V. Abgaryan, D. Alvarez-Castillo, A. Ayriyan, D. Blaschke, H. Grigorian, Universe, 4, 94 \(2018\).](#)

- ★ Surface tension effect leads to existence of pasta phases.
- ★ A parabolic interpolation method used to construct the mix phase.

$$p(\mu) = \begin{cases} p^H(\mu), & \mu \leq \mu_{cH}, \\ P^M(\mu) = \alpha_2(\mu - \mu_c)^2 + \alpha_1(\mu - \mu_c) + P_c + \Delta P, & \mu_{cH} \leq \mu \leq \mu_{cQ}, \\ p^Q(\mu), & \mu \geq \mu_{cQ} \end{cases}$$

$\alpha_1, \alpha_2, \mu_{cH}, \mu_{cQ}$ → Determined from the continuity of pressure and its derivative.

- ★ Mix Phase is parametrized by $\Delta p = \Delta P/P_c$.
- ★ $\Delta p = 0$: Maxwell Construction.



ACB4 Parametrization:

- [D. E. Alvarez-Castillo, D. Blaschke, PRC, 96, 045809, \(2017\).](#)
- [V. Paschalidis, K. Yagi, D. Alvarez-Castillo, D. Blaschke, A Sedrakian, PRD, 97, 084038, \(2018\).](#)




$$P(n) = \kappa_i \left(\frac{n}{n_0} \right)^{\Gamma_i}, \quad n_i < n < n_{i+1}, \quad i = 1, \dots, 4$$

$$P(\mu) = \kappa_i \left[(\mu - m_{0,i}) \frac{\Gamma_i - 1}{\kappa_i \Gamma_i} \right]^{\frac{\Gamma_i}{\Gamma_i - 1}}$$

i	Γ_i	κ_i [MeV fm ⁻³]	n_i [fm ⁻³]	$m_{0,i}$ [MeV]
1	4.921	2.1680	0.1650	939.56
2	0.0	63.178	0.3174	939.56
3	4.00	0.5075	0.5344	1031.2
4	2.80	3.2401	0.7500	958.55



Stellar Properties



- ❖ The second and third family merge to form a single branch for $\Delta p > 4\%$.
- ❖ Precise measurement of M - R required for detection of twin star.
- ❖ The jump $\Delta\Lambda$ (if any) can be measured $\sim 15\%$ ($< 90\%$ CI) with next-generation GW detectors ([P. Landry & K. Chakravarti, arXiv:2212.09733, 2022](#)).

Solving for f -mode Characteristics:

Perturbed Metric:

$$ds_p^2 = -e^{2\Phi(r)} dt^2 + e^{2\lambda(r)} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 + h_{\mu\nu}$$

$$h_{\mu\nu} = \begin{pmatrix} r^l H e^{2\Phi} & i\omega r^{l+1} H_1 & 0 & 0 \\ i\omega r^{l+1} H_1 & r^l H e^{2\lambda} & 0 & 0 \\ 0 & 0 & r^{l+2} K & 0 \\ 0 & 0 & 0 & r^{l+2} K \sin^2 \theta \end{pmatrix} Y_m^l e^{i\omega t},$$

[\(K. S. Thorne and A. Campolattaro, 1967\)](#)

- Cowling Approximation: background metric perturbations are neglected ($h_{\mu\nu} = 0$)
- The fluid Lagrange displacement vector (ζ)

$$\zeta^i = \left(\frac{r^l}{r} e^{-\lambda} W(r), \frac{-r^l}{r^2} V(r), 0 \right) r^{-2} Y_{lm}(\theta, \phi) \exp(i\omega t)$$

Perturbations inside the star



- Solving differential equations for coupled fluid perturbation and metric functions
- Boundary conditions:
 - Finite near center.
 - Jump condition at phase transition
 - perturbation pressure vanishes at the stellar surface

Perturbations outside the star

- Set fluid variables=0 and integrate the Zerilli equations for perturbations outside the star.
- Search for complex ω , for which one have only outgoing wave solution at infinity.

[S. Detweiler and L. Lindblom, 1985](#), [F. J. Zerilli, 1970](#), [Sotani, 2001](#).

F-mode characteristics



- Sudden increase (decrease) in the frequency (damping time) observed with appearance of twin star.
- Detections of f-mode GWs from compact stars with known mass may reveal the presence of twin stars.
- Simultaneous measurement of M - f (from binary system) can be used to comment on twin stars.

Neutron star Asteroseismology




➤ Nils Andersson and Kostas D. Kokkotas, [Mon. Not. R. Astron. Soc. 299, 1059–1068. \(1998\).](#)

$$\omega_f(\text{kHz}) \approx 0.78 + 1.635 \left(\frac{\bar{M}}{\bar{R}^3} \right)^{1/2}$$

$$\frac{1}{\tau_f(\text{s})} \approx \frac{\bar{M}^3}{\bar{R}^4} \left[22.85 - 14.65 \left(\frac{\bar{M}}{\bar{R}} \right) \right]$$

$$\omega_w(\text{kHz}) \approx \frac{1}{\bar{R}} \left[20.92 - 9.14 \left(\frac{\bar{M}}{\bar{R}} \right) \right]$$

$$\frac{1}{\tau_w(\text{ms})} \approx \frac{1}{\bar{M}} \left[5.74 + 103 \left(\frac{\bar{M}}{\bar{R}} \right) - 67.45 \left(\frac{\bar{M}}{\bar{R}} \right)^2 \right]$$



Further works improved the **Asteroseismology** problem,

- O. Benhar, V. Ferrari, and L. Gualtieri, Phys. Rev. D 70, 124015 (2004).
- L. K. Tsui and P. T. Leung, Mon. Not. R. Astron. Soc. 357, 1029 (2005).
- J. L. Blázquez-Salcedo, L. M. González-Romero, and F. Navarro-Lérída, Phys. Rev. D, 89, 044006 (2014).
- G. Lioutas and N. Stergioulas, Gen. Relativ. Gravit. 50, 12 (2018).
- H. Sotani and B. Kumar, Phys. Rev. D 104, 123002 (2021).
- T. Zhao and J. M. Lattimer, PRD 106, 123002 (2022)

- URs among f-mode characteristics (f , τ_f or $\omega=2\pi f+1/\tau_f$) and NS observables.

$$f(\text{kHz}) = a_r + b_r \sqrt{\frac{M}{R^3}}$$


→

Empirical relations (EOS dependent)

$$\text{Re}(M\omega) = a_0 + a_1 \left(\frac{M}{R}\right) + a_2 \left(\frac{M}{R}\right)^2$$

$$\text{Im}(M\omega) = b_0 \left(\frac{M}{R}\right)^4 + b_1 \left(\frac{M}{R}\right)^5 + b_2 \left(\frac{M}{R}\right)^6$$




	Re($M\omega$)		Im($M\omega$)	
a_0	$-0.027 \pm 9 \times 10^{-5}$	b_0	$(9.81 \pm 0.004) \times 10^{-2}$	
a_1	0.610 ± 0.0015	b_1	$(-4.444 \pm 0.003) \times 10^{-1}$	
a_2	0.049 ± 0.002	b_2	$(4.91 \pm 0.0045) \times 10^{-1}$	



- Scaled Universal relations are more useful.
- Twin stars do not violate the URs. So the URs can be used for EoS inference.
- URs involving tidal deformability have also been examined.

Compact star observables from f-mode observations : Role of UR Uncertainty

- ❖ With the assumption that f , τ are measured precisely.
- ❖ Errors on UR results uncertainties on M - R .



- ★ The presence of the twins may be confirmed with exact measurement of f , and τ .
- ★ The unstable branch of $\Delta p = 0\%$ can be distinguished from the connecting stable branch of $\Delta p = 8\%$.
- ★ Differentiating among $\Delta p = 0\%$ and $\Delta p = 5\%$ is more challenging.


Inclusion of Observational Uncertainties

- F-mode being excited during pulsar glitches. All the energy radiated through GW.
- The burst waveform is modelled as an exponentially damped oscillation.

$$h(t) = h_0 \exp(-t/\tau_f) \sin(2\pi\nu_f t), \quad t > 0$$

([B.J. Owen,2010](#), [Ho et al. 2020](#))


$$h_0 = 4.85 \times 10^{-17} \sqrt{\frac{E_{gw}}{M_\odot c^2}} \sqrt{\frac{0.1 \text{sec} \cdot 1 \text{kpc}}{\tau_f d}} \left(\frac{1 \text{kHz}}{\nu_f} \right)$$




[Pradhan et al., Phys. Rev. C 106\(2022\), 015805](#)

F-mode GW

- B. Abbott et al., LVC, [ApJ 874 163, 2019](#);
- R. Abbott et al., LVK, [PhRvD, 104, 122004, 2021](#).
- R. Abbott, et al., LVK, [arXiv:2210.10931, 2022](#).
- R. Abbott, et al., LVK, [arXiv:2203.12038, 2022](#).
- D. Lopez et al., [PhRvD, 106, 103037, 2022](#)



[Ho et al. PRD 101, 103009 \(2020\)](#)



[B. K. Pradhan, D. Pathak, and D. Chatterjee, arxiv.2306.04626\(2023\)](#)

$$E_{gw} = E_{\text{glitch}} = 4\pi^2 I \nu^2 \left(\frac{\Delta\nu}{\nu} \right)$$


Inclusion of Observational Uncertainties

- Parameter Estimation for GW signal parameters are carried out using **Bilby**.
- Priors are kept,
 - logUniform in E_{gw} .
 - $\nu_f \in \text{U}[800, 3500]$ Hz.
 - $\tau_f \in \text{U}[0.05, 0.7]$ s.
 - We fix the distance.


$$h(t) = h_0 \exp(-t/\tau_f) \sin(2\pi\nu_f t), \quad t > 0$$

$$h_0 = 4.85 \times 10^{-17} \sqrt{\frac{E_{\text{gw}}}{M_{\odot} c^2}} \sqrt{\frac{0.1 \text{sec} \cdot 1 \text{kpc}}{\tau_f d}} \left(\frac{1 \text{kHz}}{\nu_f} \right)$$

- Frequency can be measured accurately in A+ and ET.
- Damping time can have error ~20-50% in A+ and ~5-15% in ET.
- M-R posterior is obtained using UR.
- With a 90% CI, M can be measured to ~6% in ET.
- With a 90% CI, R can be measured to ~2% in ET.
- Error on M, R are large in A+.





UR



Inclusion of Observational Uncertainties

- Glitching pulsars data taken from the [Jodrell Bank Glitch catalogue](#).
- Spin frequency, distance (d) and sky position to each pulsar are assigned from [ATNF Pulsar Catalogue](#).
- Consider few random mass configurations with an assumed EOS model.
- Then f-mode frequency, damping time, moment of inertia to pulsars from the assumed EOS model.



- The measurement of R from f-mode observation may confirm the presence of twins.
- More challenging for low mass twins. However, we have more observations at low masses.
- Differentiating the nature of Δp is more challenging.

Summary

- ❖ F-mode oscillation of hybrid stars and twin stars involving the “pasta phase” is investigated.
- ❖ We re-examined the asteroseismology problem considering the twin stars.
- ❖ For precise f-mode measurement provides suitable scenario for twin star detection.
- ❖ F-mode GW detection with next-generation GW offers a promising scenario for confirming the existence of the twin stars.
- ❖ Distinguishing the nature of hadron-quark crossover phase transition is more challenging.
- ❖ This work can be improved considering the effect of rotation and magnetic field.
- ❖ A detailed Bayesian study is in progress to constrain the pasta phase parameters from f-mode observation.
- ❖ Speculation or constrain of twin star parameters/surface tension/pasta phase from future GW measurement from binary system is in progress.

Thank you





Twin stars in binary:

$$\mathcal{M}_c = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}}$$






$$\tilde{\Lambda} = \frac{16}{13} \left[\frac{(m_1 + 12m_2)m_1^4 \Lambda_1}{(m_1 + m_2)^{1/5}} + 1 \leftrightarrow 2 \right]$$

$$C_{DT} = -\frac{1}{X_1 X_2} \left[\frac{\Lambda_1}{(m_1 \omega_1)^2} X_1^6 (155 - 147X_1) + 1 \leftrightarrow 2 \right]$$

- ❖ There is a jump in the binary parameters in presence of twin stars.
- ❖ Tidal parameter can be more useful.
- ❖ The f-mode parameters can add more information.



Additional Slides



Introduction: Neutron star and QNMs

- Non-radial QNMs raised from time varying quadrupole deformations are source of GWs.
- These deformations contain signature of underlying composition of NS and reflect in the form of observed frequencies and damping time.

Fluid displacement vector :

$$\vec{\zeta}(\vec{r}, t) = \sum_{lm} \left[\zeta_r(r) \hat{r} + \zeta_h(r) \left(\hat{\theta} \partial_\theta + \hat{\phi} \frac{1}{\sin \theta} \partial_\phi \right) \right] Y_{lm}(\theta, \phi) e^{i\omega t}$$

([K. S. Thorne, 1967](#))

Newtonian limit (T=0, B=0, no rotation)

([Gudrun Kristine Høye, 1999](#))

- Continuity: $\rho' = -\vec{\nabla} \cdot (\rho_0 \vec{\zeta})$

☐ At center : Regularity

☐ At surface:

- Euler: $\rho_0 \vec{\zeta}_{tt} = -\vec{\nabla} p' - \rho_0 \vec{\nabla} \phi' - \rho' \vec{\nabla} \phi_0$

$$p' + \frac{dp_0}{dr} \zeta_r = 0$$

- Poisson : ~~$\vec{\nabla}^2 \phi' = 4\pi G \rho'$~~ (Cowling Approximation, [T. G. Cowling, 1941](#))

- Energy: $p' + \vec{\zeta} \cdot \vec{\nabla} p_0 = \frac{\Gamma_1 p_0}{\rho_0} \left(\rho' + \vec{\zeta} \cdot \vec{\nabla} \rho_0 \right)$