

In-Medium Vector Meson Spectral Functions from FRG


Maximilian Wiest
TU Darmstadt

R.-A. Tripolt, T. Galatyuk, L. v. Smekal, J. Wambach



Dileptons from hot and dense QCD matter

- ▶ Goal: Obtain chirally consistent EM spectral functions across the QCD phase diagram
- ▶ Identify the impact of possible phase transitions and CEP on the EM rates
--> EM spectral function calculated from analytically continued FRG flow equations



Nature Phys. 15(2019) 10, 1040-1045

What is FRG?

- ▶ Functional Renormalization Group
- ▶ If you do **NOT** like equations:


FRG is a machine which takes a Lagrange density and returns a phase diagram and Spectral Functions

- ▶ If you do like equations:

FRG approaches aim to solve the Wetterich equation for the effective average action and its functional derivatives to obtain an effective potential and n-point-Functions, dealing with thermal and quantum fluctuations consistently

Wetterich equation

- ▶ Wetterich, Phys.Lett.B 301 (1993) 90-94
- ▶ Describes change of effective average action with scale k
- ▶ Simple one loop structure, but:
 - Full propagators in the loops
 - Exact equation!
 - Non-perturbative!
- ▶ Regulators cut off fluctuations below renormalization scale k

$$\partial_k \Gamma_k = \frac{1}{2} \left(\text{---} \right) - \left(\text{---} \right)$$


R. A. Tripolt, PhD thesis, 2015

Parity doublet model: Bare action Γ_k

- ▶ Model includes $\pi, \sigma, \rho, a_1, N, N^*(1535)$
- ▶ Obeys chiral symmetry
- ▶ Has a mass term thanks to „Mirror Baryon Prescription“


$$\begin{aligned} \Gamma_k = \int d^4x \left\{ & \bar{N}_1 (\not{\partial} - \mu_B \gamma_0 + h_{s,1}(\sigma + i\vec{\tau} \cdot \vec{\pi} \gamma^5) + h_{v,1}(\gamma_\mu \vec{\tau} \cdot \vec{\rho}_\mu + \gamma_\mu \gamma^5 \vec{\tau} \cdot \vec{a}_{1,\mu})) N_1 \\ & + \bar{N}_2 (\not{\partial} - \mu_B \gamma_0 + h_{s,2}(\sigma - i\vec{\tau} \cdot \vec{\pi} \gamma^5) + h_{v,2}(\gamma_\mu \vec{\tau} \cdot \vec{\rho}_\mu - \gamma_\mu \gamma^5 \vec{\tau} \cdot \vec{a}_{1,\mu})) N_2 + m_{0,N} (\bar{N}_1 \gamma^5 N_2 - \bar{N}_2 \gamma^5 N_1) \\ & + U_k(\phi^2) - c\sigma + \frac{1}{2}(D_\mu \phi)^\dagger D_\mu \phi - \frac{1}{4} \text{tr} \partial_\mu \rho_{\mu\nu} \partial_\sigma \rho_{\sigma\nu} + \frac{m_v^2}{8} \text{tr} \rho_{\mu\nu} \rho_{\mu\nu} \right\} + \Delta \Gamma_{\pi a_1}. \end{aligned}$$

$m_{0,N}$ [MeV]	$h_{s,1}$ $= h_{v,1}$	$h_{s,2}$ $= h_{v,2}$	$f_\pi \equiv \sigma_0$ [MeV]	m_π [MeV]	m_σ [MeV]	m_{N_1} [MeV]	m_{N_2} [MeV]
800	6.94073	13.3493	92.8	137	474	938	1533

Tripolt et al., Phys.Rev.D 104 (2021) 5, 054005

Effective potential


- ▶ Insert bare action Γ_Λ into Wetterich equation, get flow equation for U_k
- ▶ Potential has a minimum somewhere in field configuration at $k=0$.
- ▶ Position of minimum shows breaking of underlying symmetries
 - Order parameter!
 - Phase diagram!




Lancaster, Blundell,
QFT for gifted amateur

Phase diagram in parity doublet model

Phase diagram in parity doublet model



Mass degeneracy



Tripolt et al., Phys.Rev.D 104 (2021) 5, 054005

Consequences of mirror prescription

- ▶ Remember:

$$m_{0,N}(\bar{N}_1\gamma^5 N_2 - \bar{N}_2\gamma^5 N_1)$$

- ▶ In mirror prescription, chiral symmetry breaking explains part of baryon mass generation, but not full picture
- ▶ Instead, additional mass generated by different source
 - E.g. QCD scale anomaly, see e.g. [Shifman et al., Phys.Lett.B 78 \(1978\) 443-446](#)
- ▶ Possible experimental signals for mirror prescription?
 - Hadronic signals (η -meson enhancement)? [Larionov, v. Smekal, Phys.Rev.C 105 \(2022\)](#)
- ▶ Other signals?

Prescription shows up in ρ spectral function!


$$\partial_k \Gamma_{\rho, k}^{(2)} = \text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3} - 2 \text{Diagram 4} - \frac{1}{2} \text{Diagram 5}$$

The equation shows the expansion of the functional derivative of the 2-point function $\partial_k \Gamma_{\rho, k}^{(2)}$. The terms are represented by Feynman-like diagrams. The first three terms involve ρ fields and either π or a_1 fields. The fourth term involves ρ and N fields. The fifth term involves π and ρ fields. The fourth term is highlighted with a red box.

- ▶ Functional derivatives of Γ_k w.r.t. fields give n-point functions
- ▶ Mirror prescription gives rise to low energy peak in ρ $\Gamma^{(2)}$ function due to $\rho + N_1 \rightarrow N_2$
 - Unique to parity doublet model with mirror prescription!

Tripolt et al., Phys.Rev.D 104 (2021) 5, 054005

Example: $T=40$ & $\mu_B=890$, $p=0$ MeV, $\text{Im } \Gamma_\rho^2$



Prescription shows up in ρ spectral function!


$$\partial_k \Gamma_{\rho, k}^{(2)} = \text{Diagram 1} + \text{Diagram 2} + \text{Diagram 3} - 2 \text{Diagram 4} - \frac{1}{2} \text{Diagram 5}$$

- ▶ Functional derivatives of Γ_k w.r.t. fields give n-point functions
- ▶ Mirror prescription gives rise to low energy peak in ρ $\Gamma^{(2)}$ function due to $\rho + N_1 \rightarrow N_2$
 - Unique to parity doublet model with mirror prescription!
- ▶ Also manifests in spectral function!

$$\Pi_\rho \propto \frac{\text{Im } \Gamma_\rho^{(2)}}{\left(\text{Re } \Gamma_\rho^{(2)}\right)^2 + \left(\text{Im } \Gamma_\rho^{(2)}\right)^2}$$

Tripolt et al., Phys.Rev.D 104 (2021) 5, 054005

Example: $T=40$ & $\mu_B=890$, $p=0$ MeV



Spectral function to dilepton spectra

- ▶ Important equation for extraction of thermal dilepton spectra


$$\frac{dN_{ll}}{d^4x d^4q} = -\frac{\alpha_{EM}^2}{\pi^3 M^2} L(M^2) f^{BE}(q_0, T) \text{Im}\Pi_{EM}(M, q, \mu_B, T).$$

- ▶ Vector Meson Dominance: Vector SF proportional to electromagnetic SF
- ▶ Peak translates from spectral function to dilepton spectra!

Takeaway: Dilepton yield depends on T, μ_B (ρ_B), is obtained by integrating over space-time and 4-momentum

McLerran, Toimela, Phys. Rev. D 31 (1985), p. 545


Mirror prescription in dilepton yield



- ▶ Take ρ, T distributions from UrQMD via Coarse Graining procedure
- ▶ For $\vec{p}=0 \text{ MeV}$ spectral function, peak is seen clearly in FRG SF
- ▶ Comparison with Rapp-Wambach spectral function
 - Describes spectra over wide range of energies

R. Rapp, J. Wambach, Eur. Phys. J. A 6, 415 (1999)

Mirror prescription in dilepton yield




► Low energies?

- 1st step: Full momentum dependence!
- Does the peak survive for finite momentum?


$$\rho = \frac{1}{3}(2\rho_{\perp} + \rho_{||})$$

R. Rapp, J. Wambach, Eur. Phys. J. A 6, 415 (1999)


Example: $T=40$ & $\mu_B=890$, $p=0$ MeV




Example: $T=40$ & $\mu_B=890$, $p=50$ MeV




Example: $T=40$ & $\mu_B=890$, $p=100$ MeV




Example: $T=40$ & $\mu_B=890$, $p=150$ MeV




Example: $T=40$ & $\mu_B=890$, $p=200$ MeV




Example: $T=40$ & $\mu_B=890$, $p=250$ MeV




Example: $T=40$ & $\mu_B=890$, $p=300$ MeV




Example: $T=40$ & $\mu_B=890$, $p=350$ MeV




Example: $T=40$ & $\mu_B=890$, $p=400$ MeV




Example: $T=40$ & $\mu_B=890$, $p=450$ MeV




Example: $T=40$ & $\mu_B=890$, $p=500$ MeV




Example: $T=40$ & $\mu_B=890$, $p=550$ MeV




Example: $T=40$ & $\mu_B=890$, $p=600$ MeV




Example: $T=40$ & $\mu_B=890$, $p=650$ MeV




Example: $T=40$ & $\mu_B=890$, $p=700$ MeV




Example: $T=40$ & $\mu_B=890$, $p=750$ MeV




Example: $T=40$ & $\mu_B=890$, $p=800$ MeV




Example: $T=40$ & $\mu_B=890$, $p=850$ MeV




Example: $T=40$ & $\mu_B=890$, $p=900$ MeV




Example: $T=40$ & $\mu_B=890$, $p=950$ MeV




Example: $T=40$ & $\mu_B=890$, $p=1000$ MeV



Example: $T=40$ & $\mu_B=890$, $p=1050$ MeV




Dilepton invariant mass spectrum for $p>0$



- ▶ Integrated yield rises by a factor of ~ 2
- ▶ Peak is smeared out considerably

R. Rapp, J. Wambach, Eur. Phys. J. A 6, 415 (1999)

Dilepton invariant mass spectrum for $p>0$



- ▶ Integrated yield rises by a factor of ~ 2
- ▶ Peak is smeared out considerably
- ▶ Still many things missing for “realistic” spectral function
 - More baryon species
 - Pion modifications


R. Rapp, J. Wambach, Eur. Phys. J. A 6, 415 (1999)

Polarisation?

- ▶ Polarisation of thermal virtual photons (in helicity frame) given by

$$\lambda_\theta = \frac{\rho_\perp - \rho_{||}}{\rho_\perp + \rho_{||}}$$



- ▶ λ_θ describes anisotropy of decay distribution
- ▶ Can give information about origin of thermal dileptons





Speranza et al., Phys.Lett.B 782 (2018) 395-400
Seck et al., arXiv:2309.03189

Faccioli, Lourenco, Lect.Notes Phys. 1002 (2022)



Example: $T=40$ & $\mu_B=890$, $p=0$ MeV





Example: $T=40$ & $\mu_B=890$, $p=50$ MeV





Example: $T=40$ & $\mu_B=890$, $p=100$ MeV





Example: $T=40$ & $\mu_B=890$, $p=150$ MeV





Example: $T=40$ & $\mu_B=890$, $p=200$ MeV





Example: $T=40$ & $\mu_B=890$, $p=250$ MeV





Example: $T=40$ & $\mu_B=890$, $p=300$ MeV





Example: $T=40$ & $\mu_B=890$, $p=350$ MeV





Example: $T=40$ & $\mu_B=890$, $p=400$ MeV





Example: $T=40$ & $\mu_B=890$, $p=450$ MeV





Example: $T=40$ & $\mu_B=890$, $p=500$ MeV





Example: $T=40$ & $\mu_B=890$, $p=550$ MeV





Example: $T=40$ & $\mu_B=890$, $p=600$ MeV





Example: $T=40$ & $\mu_B=890$, $p=650$ MeV





Example: $T=40$ & $\mu_B=890$, $p=700$ MeV





Example: $T=40$ & $\mu_B=890$, $p=750$ MeV





Example: $T=40$ & $\mu_B=890$, $p=800$ MeV





Example: $T=40$ & $\mu_B=890$, $p=850$ MeV





Example: $T=40$ & $\mu_B=890$, $p=900$ MeV





Example: $T=40$ & $\mu_B=890$, $p=950$ MeV



Example: $T=40$ & $\mu_B=890$, $p=1000$ MeV




Example: $T=40$ & $\mu_B=890$, $p=1050$ MeV




Polarisation: Comparison to Rapp-Wambach


This Work



Rapp-Wambach



Medium Effects: Transverse vs. Longitudinal



M. Urban, PhD-Thesis, 2001

Summary and Outlook


- ▶ Dilepton invariant mass spectra could, at low momenta, provide information on mirror baryon scenario
- ▶ Finite momentum modifies low energy limit strongly, increases dilepton production
 - Peak structure is washed out integrating over momenta
- ▶ Dilepton polarization could provide additional insights
 - Provided the rich structure obtained in the PDM survives the inclusion of additional effects
- ▶ More baryons, medium modifications of pions to be included!

Appendix


Spectral function

- ▶ More precisely: An electromagnetic spectral function
 - Even more precisely: A vector-meson spectral function
 - Which is equivalent up to a constant in the Vector Dominance Model

- ▶ Mainly two:
 - Hadronic many body approach by Rapp
 - FRG approach by Tripolt



Rapp, Physics Letters B, Volume 731, 2014, Pages 103-109



Tripolt, Phys.Rev.D 104 (2021)

Guide to deriving the Wetterich Equation


► Legendre-Transform of $\log Z[J]$ gives generating functional of 1-PI-Diagrams

- effective action Γ
- 1PI: Diagrams, which cannot be cut in 2 parts by cutting 1 line

► Add a fictional mass term $\propto m_k^2 \phi^2$ to Lagrangian

$$Z[J] = \int D\phi \exp(-\int d^4x \mathcal{L}[\phi(x)] + J(x)\phi(x))$$


$$Z_k[J] = \int D\phi \exp(-\int d^4x \mathcal{L}[\phi(x)] + J(x)\phi(x) + m_k^2\phi(x)^2)$$



Lancaster, Blundell,
QFT for gifted amateur

NOT 1-PI:


► Mass term m_k^2 often called „Regulator“ and written R_k



Guide to deriving the Wetterich Equation

- ▶ Mass term cuts away fluctuations with momentum scale $< k$
- ▶ Define Γ_k
 - Same as before, but averages action over Volume $1/k^3$
 - Call Γ_k effective AVERAGE action
- ▶ Find change $\partial_k \Gamma_k$ with k
- ▶ Described by Wetterich equation
 - follows from very little assumptions of form of R_k
 - Uses in (but not limited to) QCD, magnets, condensed matter, statistical physics, critical phenomena

Quick Rundown:

$$\partial_k \Gamma_k = \frac{1}{2} \left(\text{---} \right) - \text{---}$$



- ▶ Lines: (Euclidean) propagators
- ▶ Circles: Regulators $\propto \theta(k^2 - q^2)$
- ▶ Trace to be taken over all internal degrees of freedom: Isospin space, parity indices, fermion indices, Lorenz indices, internal momenta, etc.

Parity in linear σ model


► Inclusion of parity partners in linear σ model:

- 2 parity conserving ways:

„naive“ prescription



„mirror“ prescription



Consequences of prescriptions

- ▶ In both cases, mass terms for single baryon species are not allowed



$$\mathcal{L}_m = m\bar{\psi}_i\psi_i$$

- ▶ However: In mirror case, mixing allows for mass term

$$\mathcal{L}_m = m(\bar{\psi}_2\psi_1 + \bar{\psi}_1\psi_2)$$

- ▶ Chirally restored phase: Massless baryons vs. degenerate massive baryons

Weyrich et al., Phys.Rev.C 92 (2015) 1, 015214