

# QCD

## problem set 6

1. In this problem we will discuss parton properties assuming very simple models for quark and gluon distributions. My advice is to use *Mathematica* for these calculations.

- (a) Using properties of parton distributions given at lecture 2 in section *Quarks as Partons* calculate normalization constant  $A$  assuming

$$u_v(x) = \frac{2A}{\sqrt{x}}(1-x)^3, \quad d_v(x) = \frac{A}{\sqrt{x}}(1-x)^3$$

where index  $v$  stands for *valence*. Recall that total  $u$  or  $d$  quark distribution is given as a sum of valence quarks and sea quarks  $u_s$  or  $d_s$  respectively. We assume that sea quark distributions are equal to antiquark distributions

$$u_s(x) = \bar{u}(x), \quad d_s(x) = \bar{d}(x).$$

For this problem we assume that there are no strange quarks in the nucleon and we assume isospin symmetry, which says that  $u$  and  $d$  distributions in neutron, are equal to  $d$  and  $u$  distributions in proton. Calculate  $A$ . Check the value of charge of the proton and neutron. Calculate total momentum carried by the valence quarks. At this point we do not need any information on the sea quarks.

- (b) Gottfried sum rule. Calculate the difference of the structure functions of the proton and neutron:

$$S_G = \int_0^1 \frac{dx}{x} (F_2^p(x) - F_2^n(x)).$$

Experimental value reads  $S_G \simeq 0.24$ . As you will see  $S_G$  will depend on the integral over the distributions of the sea quarks. Assume the sea quark distribution of the following form

$$\bar{u}(x) = \frac{B}{x}(1-x)^8, \quad \bar{d}(x) = \frac{B}{x}(1-x)^\beta.$$

Note that constant  $B$  must be the same in both cases to assure that  $S_G$  is finite. From the experimental value of  $S_G$  calculate  $B$  for several choices of power  $\beta$  taking a few values around 8. Note that the antiquark and sea distributions must be positive. For these choices calculate total momentum carried by quarks. Is it possible to get the value of 100%?

- (c) Choose gluon distribution of the form

$$g(x) = Cx(1-x)^4.$$

For given  $\beta$  from the previous problem calculate  $C$  from the condition that the total momentum of the proton is 1.