

# Lorentzian spectral $\zeta$ -functions

joint work with [Nguyen Viet Dang](#) (Jussieu)



“Noncommutative geometry: metric and spectral aspects”, Kraków, 2022

[Michał Wrochna](#)  
Cergy Paris Université

*based on joint works:*

---

w. Nguyen Viet Dang ([Jussieu](#))

[arXiv:2012.00712](#), *J. Eur. Math. Soc. (JEMS)*

[arXiv:2108.07529](#), *J. Éc. polytech. Math.*

[arXiv:2202.06408](#), proceedings paper

---

w. Ruben Zeitoun ([Cergy & ENS Lyon](#))

[arXiv:2204.01094](#), in: *Analysis and Partial Differential Equations, Trends in Mathematics, Springer*

---

w. Nguyen Viet Dang ([Jussieu](#)) & András Vasy ([Stanford](#))

 [work in progress](#)

---

# Introduction

Consider a **Lorentzian manifold**  $(M, g)$ .

The metric has signature  $(+, -, \dots, -)$ .

For instance Minkowski space:  $\mathbb{R}^{1+d}$ ,  $g_0 = dt^2 - dy_1^2 - \dots - dy_{n-1}^2$ .

The Lorentzian **Laplace–Beltrami operator** or **wave operator**:

$$\square_g = \sum_{i,j=0}^{n-1} |g(x)|^{-\frac{1}{2}} \partial_{x^i} |g(x)|^{\frac{1}{2}} g^{ij}(x) \partial_{x^j}$$

on Minkowski,  $\square_g = \partial_t^2 - (\partial_{y_1}^2 + \dots + \partial_{y_{n-1}}^2)$

$\square_g$  (+ non-linearity) has rich theory of solving **Cauchy problem**, **asymptotic analysis of solutions**, **propagation of singularities**, etc.

*Relatively recently:* **global** theory of  $\square_g$  (Fredholm property, Hilbert space invertibility) **Vasy '13 et al.**. Techniques of **microlocal** and **asymptotic analysis** in relation with classical dynamics and geometry.

As opposed to  $\Delta_g$  on Riemannian manifold,  $\square_g$  is **non-elliptic**.

Recently established **non-elliptic Fredholm theory** for problems such as :

1. stability of black hole solutions of Einstein equations
2. Anosov/Morse–Smale flows, dynamical zeta functions
3. Quantum Field Theory on curved spacetimes  
(Vasy, Gérard–W., Nakamura–Taira, Dang–W., Bär–Strohmaier, ...)

striking similarities with Euclidean setting  $\Delta_g$ ! (inverses, essential self-adjointness)

? How is **global**  $\square_g$  related to **geometry** of  $(M, g)$ ?

(+ related questions at the interface of quantum physics, gravity and NCG)

# Spectral zeta function

$(M, g)$  compact Riemannian  $\implies \Delta_g$  has discrete spectrum.

Recall Riemann zeta  $\zeta(\alpha) = \sum_{\lambda=1}^{\infty} \lambda^{-\alpha}$ , then **spectral zeta**:

$$\mathbb{C} \ni \alpha \mapsto \zeta_{\Delta}(\alpha) = \sum_{\lambda \in \text{sp}(\Delta_g) \setminus \{0\}} \lambda^{-\alpha}.$$

**Theorem** (Minakshisundaram–Pleijel, Seeley)

*The function  $\zeta_{\Delta}(\alpha) = \text{Tr}_{L^2}(\Delta_g^{-\alpha})$  is **holomorphic** on  $\text{Re } \alpha > \frac{n}{2}$ , with **meromorphic continuation** to  $\alpha \in \mathbb{C}$  and poles at  $\{\frac{n}{2}, \frac{n}{2} - 1, \dots, 1\}$ .*

# Spectral zeta function

$(M, g)$  compact Riemannian  $\implies \Delta_g$  has discrete spectrum.

Recall Riemann zeta  $\zeta(\alpha) = \sum_{\lambda=1}^{\infty} \lambda^{-\alpha}$ , then **spectral zeta**:

$$\mathbb{C} \ni \alpha \mapsto \zeta_{\Delta}(\alpha) = \sum_{\lambda \in \text{sp}(\Delta_g) \setminus \{0\}} \lambda^{-\alpha}.$$

## Theorem (Minakshisundaram–Pleijel, Seeley)

The function  $\zeta_{\Delta}(\alpha) = \text{Tr}_{L^2}(\Delta_g^{-\alpha})$  is **holomorphic** on  $\text{Re } \alpha > \frac{n}{2}$ , with **meromorphic continuation** to  $\alpha \in \mathbb{C}$  and poles at  $\{\frac{n}{2}, \frac{n}{2} - 1, \dots, 1\}$ .

+local version with densities:

$\alpha \mapsto \Delta_g^{-\alpha}(x, x)$  **holomorphic** on  $\text{Re } \alpha > \frac{n}{2}$ , with **meromorphic continuation** to  $\alpha \in \mathbb{C}$  and poles at  $\{\frac{n}{2}, \frac{n}{2} - 1, \dots, 1\}$ , **smooth** in  $x \in M$ .

Here  $\Delta_g^{-\alpha}(x, x')$  is the **Schwartz kernel** of  $\Delta^{-\alpha}$ , so

$$\text{Tr}_{L^2}(\Delta_g^{-\alpha}) = \int_M \Delta_g^{-\alpha}(x, x) dx$$

# The spectral action principle

The **heat kernel expansion** (small  $t$  expansion of  $e^{-t\Delta_g}(x, x)$ ) relates  $\Delta_g$  with invariants, in particular **scalar curvature**  $R_g(x)$ .

**Theorem** (elliptic theory + Connes, Kalau–Walze, Kastler)

When  $\dim(M) = n \geq 4$ ,

$$\operatorname{res}_{\alpha=\frac{n}{2}-1} \operatorname{Tr}_{L^2} (\Delta_g^{-\alpha}) = \frac{\int_M R_g(x)}{6(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} - 1)}.$$

*Local version for diagonal value  $x = x'$  of Schwartz kernel  $\Delta_g^{-\alpha}(x, x')$ :*

$$\operatorname{res}_{\alpha=\frac{n}{2}-1} \Delta_g^{-\alpha}(x, x) = \frac{R_g(x)}{6(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} - 1)}.$$

- This is a **spectral action** for Euclidean gravity:  $\delta_g R_g = 0$  is equivalent to **Einstein equations**.
- Poles are geometric  $\Rightarrow$  locality of counterterms in **zeta function regularisation** in QFT **Hawking '77**
- The analytic residue equals a **Guillemin–Wodzicki residue** (or **non-commutative residue**), therefore a Dixmier trace **Connes '88**

## Theorem (elliptic theory + Chamseddine–Connes)

For any Schwartz function  $f$ ,

$$f(\Delta_g/\lambda^2)(x, x) = \sum_{j=0}^N \lambda^{n-2j} C_j(f) a_j(x) + \mathcal{O}(\lambda^{n-2N-1}),$$

where  $a_j(x)$  are the heat kernel coefficients.

- Vector bundle version useful for **Dirac**  $f\left(\frac{D^2}{\lambda^2}\right)$ .
- Twisting the bundle yields **Standard model** Lagrangian.



## Theorem (elliptic theory + Chamseddine–Connes)

For any Schwartz function  $f$ ,

$$f(\Delta_g/\lambda^2)(x, x) = \sum_{j=0}^N \lambda^{n-2j} C_j(f) a_j(x) + \mathcal{O}(\lambda^{n-2N-1}),$$

where  $a_j(x)$  are the heat kernel coefficients.

- Vector bundle version useful for **Dirac**  $f\left(\frac{D^2}{\lambda^2}\right)$ .
  - Twisting the bundle yields **Standard model** Lagrangian.
- 

But **no direct physical meaning** unless  $(M, g)$  Lorentzian...

Yet, fundamental difficulties: Lorentzian  $\square_g$  **not elliptic**, **not bounded from below**. There is **no Lorentzian heat kernel**.

# Lorentzian non-commutative geometry, a work in progress



Spectral triples by Wick rotation or spacetime foliations

van den Dungen–Paschke–Rennie '13, van den Dungen–Rennie '16, van den Dungen '18 ...



Krein space based spectral triples

Suijlekom '04, Strohmaier '06, Paschke–Sitarz '06, Barrett '07, Besnard '16, van den Dungen '16 ...



Lorentzian distance function and causal relations

Moretti '03, Besnard '09, Franco '10–'18, Rennie–Whale '17, Minguzzi '17, Franco–Eckstein '13–'15, Bizi–Besnard '17 ...



Algebraic structure of Lorentzian spectral triples / actions

Franco '12, Bochniak–Sitarz '18, Bizi–Brouder–Besnard '18 ...



Wick rotation of the spectral action

D'Andrea–Kurkov–Lizzi '16, Devastato–Farnsworth–Lizzi–Martinetti '18, Martinetti–Singh '19 ...

# Is there a spectral action with $\square_g$ ?

For  $(M, g)$  Lorentzian,  $\Delta_g$  becomes  $\square_g$ . Two hints:

1. The **local geometric quantities** (e.g.  $R_g(x)$ ) still make sense.
  - Lorentzian version of local **heat kernel coefficients**  $a_j(x)$  by solving analogous transport equations
  - formal **Hadamard parametrix** for  $\square_g$  produces  $a_j(x)$
2. Recent results show **essential self-adjointness** of  $\square_g$ :
  - Static spacetimes (e.g.  $\partial_t^2 - \Delta_h$  with **time-independent coefficients**): Dereziński-Siemssen '18
  - For perturbations of Minkowski space (and more general **non-trapping Lorentzian scattering spaces**):  
Vasy '20, Nakamura-Taira '20  
(related results: Gérard-Wrochna '19-'20, Kamiński '19,  
Dereziński-Siemssen '19, Colin de Verdière-Le Bihan '20, Taira '20)
  - Classes of **asymptotically static spacetimes** Nakamura-Taira '22

$\Rightarrow f(\square_g)$  well-defined!

But is there any relationship between **1.** and **2.** like in elliptic case?

## **I. Main results**

# Main theorem

Assume  $(M, g)$  is a perturbation of Minkowski space (or more general non-trapping Lorentzian scattering space, see later), of even dimension  $n$ .

## Theorem (Dang, Wrochna '20)

For  $\varepsilon > 0$ , the Schwartz kernel of  $(\square_g - i\varepsilon)^{-\alpha}$  has for  $\operatorname{Re} \alpha > \frac{n}{2}$  a well-defined on-diagonal restriction  $(\square_g - i\varepsilon)^{-\alpha}(x, x)$ , which extends as a meromorphic function of  $\alpha \in \mathbb{C}$  with poles at  $\{\frac{n}{2}, \frac{n}{2} - 1, \frac{n}{2} - 2, \dots, 1\}$ . Furthermore,

$$\lim_{\varepsilon \rightarrow 0^+} \operatorname{res}_{\alpha = \frac{n}{2} - 1} (\square_g - i\varepsilon)^{-\alpha}(x, x) = \frac{R_g(x)}{i6(4\pi)^{\frac{n}{2}} \Gamma(\frac{n}{2} - 1)},$$

where  $R_g(x)$  is the scalar curvature at  $x \in M$ .

- **Spectral action for gravity!** Proof directly in Lorentzian signature. Perturbations of Minkowski included (no symmetries assumed).
- The  $\varepsilon \rightarrow 0^+$  avoids low-frequency problems and responsible for relationship with Feynman propagator.

## Main theorem 2

**Theorem** (Dang, Wrochna '20)

For any Schwartz  $f$  with Fourier transform in  $]0, +\infty[$ ,

$$f((\square_g + i\varepsilon)/\lambda^2)(x, x) = \sum_{j=0}^N \lambda^{n-2j} C_j(f) a_j(x) + \mathcal{O}(\varepsilon, \lambda^{n-2N-1}),$$

where  $a_j(x)$  are Hadamard coefficients.

**Theorem** (Dang, Wrochna '21)

Poles of  $\zeta_{g,\varepsilon}(\alpha) := (\square_g - i\varepsilon)^{-\alpha}(x, x)$  can be recovered by a scaling procedure.

(This generalizes the *Guillemin–Wodzicki residue* of  $\Psi$ DOs.)

# Scaling towards the diagonal

Let  $\Delta = \{(x, x) \mid x \in M\}$ .

A vector field  $X$  is **radial** (or **Euler**) if  $Xf = f$  modulo quadratically vanishing terms for all  $f$  with  $f|_{\Delta} = 0$ .

Locally there are coordinates  $(x^i, h^i)_{i=1}^n$  s.t.  $\Delta = \{h^i = 0\}$  and  $X = \sum_{i=1}^n h^i \partial_{h^i}$ .

$u \in \mathcal{D}'_{\Gamma}(\mathcal{U})$  is **log-polyhomogeneous** if

$$e^{-tX}u = \sum_{p \leq k \leq N, 0 \leq i \leq l-1} e^{-tk} \frac{(-1)^i t^i}{i!} (X - k)^i u_k + \mathcal{O}_{\mathcal{D}'_{\Gamma}(\mathcal{U})}(e^{-t(N+1-\varepsilon)}).$$

**Pollicott–Ruelle resonances** of the flow  $e^{-tX}$  are the poles of

$$\int_0^{\infty} e^{-tz} \langle (e^{-tX}u), \varphi \rangle dt = \sum_{k=p, 0 \leq i \leq l-1}^N (-1)^i \frac{\langle (X - k)^i u_k, \varphi \rangle}{(z + k)^{i+1}}$$

+ holomorphic on  $\operatorname{Re} z \leq N$ .

# Dynamical definition of residue

Suppose  $\Gamma|_{\Delta} \subset N^* \Delta$ . Let  $\Pi_0 :=$  projection on zero resonance.

The **dynamical residue of  $\mathcal{K}$**  (w.r.t.  $X$ ) is:

$$\text{res}_X \mathcal{K} = \iota_{\Delta}^* (X(\Pi_0(\mathcal{K}))) \in C^{\infty}(M).$$

Might be ill-defined, and might depend on  $X$ . But...

## Theorem (Dang–Wrochna '21)

For all radial  $X$  and all  $k = 1, \dots, \frac{n}{2}$  and  $\varepsilon > 0$ ,

$$\text{res}_{\alpha=k} \zeta_{g,\varepsilon}(\alpha) = \frac{1}{2} \text{res}_X \left( (\square_g - i\varepsilon)^{-k} \right),$$

where  $\zeta_{g,\varepsilon}(\alpha)$  is the **spectral zeta function density** of  $\square_g - i\varepsilon$ .


“Analytic residues of  $\zeta_{g,\varepsilon}$  are **dynamical residues** (scaling anomalies).”



## II. From resolvent of $\square_g$ to geometric invariants

# General plan of proof

- 1) Let  $P = \square_g$  on Lorentzian  $(M, g)$ . If resolvent exists  $(P - i\varepsilon)^{-\alpha}$  as contour integral of  $(P - z)^{-1}$ . For  $\alpha = N + \mu > 0$ :



$$(P - i\varepsilon)^{-\alpha} = \frac{1}{2\pi i} \int_{\gamma_\varepsilon} (z - i\varepsilon)^{-\mu} (P - i\varepsilon)^{-N} (P - z)^{-1} dz, \quad .$$

- 2) Construct a **Hadamard parametrix**  $H_N(z)$  (replaces heat kernel) and **show it approximates the resolvent** uniformly in  $z$ .
- 3) Deduce **regularity properties**, compute **poles** and get **curvature**  $R$  from contour integrals of  $H_N(z)$ .

### Construction of Hadamard parametrix $H_N(z)$ :

Let  $\mathbf{F}_\alpha(z, |\cdot|_g)$  be locally given by

$$F_\alpha(z, x) = \frac{1}{\Gamma(\alpha + 1)(2\pi)^n} \int e^{i\langle x, \xi \rangle} (|\xi|_{g_0}^2 - i0 - z)^{-\alpha-1} d^n \xi$$

(in normal coordinates) then ansatz of order  $N$ :

$$H_N(z, \cdot) = \sum_{k=0}^N u_k \mathbf{F}_k(z, |\cdot|_g) \in \mathcal{D}'(\mathcal{U}).$$

solved modulo errors by transport equations thanks to

$$(P - z)(u \mathbf{F}_\alpha) = \alpha u \mathbf{F}_{\alpha-1} + (Pu) \mathbf{F}_\alpha + (hu + 2\rho u) \frac{\mathbf{F}_{\alpha-1}}{2}$$

for all  $u \in C^\infty(M)$ , where  $h(x) = b^j(x)g_{0,jk}x^k$  and  $\rho = x^k \partial_{x^k}$ .

## Compute poles and get curvature:

Now  $(P - i\varepsilon)^{-\alpha}(x, x)$  expressed by contour integrals of  $\mathbf{F}_\beta(z, \cdot)$ .

$$\frac{1}{2\pi i} \int_{\gamma_\varepsilon} (z - i\varepsilon)^{-\alpha} \mathbf{F}_k(z, \cdot) dz = \frac{(-1)^k \Gamma(-\alpha + 1)}{\Gamma(-\alpha - k + 1) \Gamma(\alpha + k)} \mathbf{F}_{k+\alpha-1}(i\varepsilon, \cdot)$$

Residue computation by **homological argument**.

scalar curvature in normal coordinates comes from

$$P = \partial_{x^k} g^{kj}(x) \partial_{x^j} + g^{jk}(x) (\partial_{x^j} \log |g(x)|^{\frac{1}{2}}) \partial_{x^k},$$

transport equation  $u_1(0) = -P u_0(0) = -P(|g(0)|^{\frac{1}{4}} |g(x)|^{-\frac{1}{4}})|_{x=0}$   
and  $g_{ij}(x) = g_{0,ij} + \frac{1}{3} R_{ikjl} x^k x^l + \mathcal{O}(|x|^3)$ .

## Hadamard parametrix $H_N$ approximates $(P - z)^{-1}$ ?

$$(P - z) \left( \sum_{k=0}^N u_k \mathbf{F}_k(z, \cdot) \chi \right) = |g|^{-\frac{1}{2}} \delta_\Delta + (Pu_N) \mathbf{F}_N(z, \cdot) \chi + r_N(z),$$

where  $Pu_N$  highly regular, and  $r_N$  singular (but 0 near diagonal).  
Applying  $(P - z)^{-1}$  well-defined and yields good errors if  $(P - z)^{-1}$  is shown to have special structure of singularities and mapping properties uniformly in  $z$ .

Think of the distribution  $(x - i0)^{-1}$  on  $\mathbb{R}$ : it is singular at  $x = 0$ , but has good multiplicative properties like  $(x - i0)^{-1}(x - i0)^{-1} = (x - i0)^{-2}$ .

Here, “controlling singularities” means showing existence of  $B_1, B_2 \in \Psi^0(M)$ , as elliptic as possible s.t.

$$B_1(P - z)^{-1} B_2^* : L^2(M) \rightarrow C^\infty(M)$$

with seminorms  $O(1 + |z|)^{-m}$ .

Related problems in QFT: singularities of two-point functions  $\langle \Omega | \phi(x) \phi(x') \Omega \rangle$  as  $x \rightarrow x'$  in relationship with spacetime geometry.

### **III. Analysis of $(P - z)^{-1}$**

Suppose  $\square_g = \partial_t^2 - \Delta$ ,  $\text{Im } z > 0$ . Retarded propagator of  $P - z$ :


$$\theta(t-s) \frac{e^{i(t-s)\sqrt{-\Delta-z}} - e^{-i(t-s)\sqrt{-\Delta-z}}}{2i\sqrt{-\Delta-z}}$$

Looks like **no chance of**  $\|(\square_g - z)^{-1}\| \leq |\text{Im } z|^{-1}$ . But:

Suppose  $\square_g = \partial_t^2 - \Delta$ ,  $\text{Im } z > 0$ . Retarded propagator of  $P - z$ :

$$\theta(t-s) \frac{e^{i(t-s)\sqrt{-\Delta-z}} - e^{-i(t-s)\sqrt{-\Delta-z}}}{2i\sqrt{-\Delta-z}}$$

Looks like **no chance of**  $\|(\square_g - z)^{-1}\| \leq |\text{Im } z|^{-1}$ . But:



“Every particle in Nature has an amplitude to move backwards in time, and therefore has an anti-particle.”

– Richard Feynman

$$((\square_g - z)^{-1}u)(t, \cdot) = -\frac{1}{2} \int \frac{e^{-i|t-s|\sqrt{-\Delta-z}}}{\sqrt{-\Delta-z}} u(s, \cdot) ds. \quad (1)$$

The boundary value  $(\square_g - i0)^{-1}$  is the **Feynman propagator**.

But for general  $\square_g$  with  $t$ -dependent coefficients, nothing like (1) exists...


💡 Start with (1) at **infinity**, then propagate!



Suppose  $\square_g = \partial_t^2 - \Delta$ ,  $\text{Im } z > 0$ . Retarded propagator of  $P - z$ :

$$\theta(t-s) \frac{e^{i(t-s)\sqrt{-\Delta-z}} - e^{-i(t-s)\sqrt{-\Delta-z}}}{2i\sqrt{-\Delta-z}}$$

Looks like **no chance of**  $\|(\square_g - z)^{-1}\| \leq |\text{Im } z|^{-1}$ . But:



“Every particle in Nature has an amplitude to move backwards in time, and therefore has an anti-particle.”

– Richard Feynman

$$((\square_g - z)^{-1}u)(t, \cdot) = -\frac{1}{2} \int \frac{e^{-i|t-s|\sqrt{-\Delta-z}}}{\sqrt{-\Delta-z}} u(s, \cdot) ds. \quad (1)$$

The boundary value  $(\square_g - i0)^{-1}$  is the Feynman propagator.



Use **radial estimates** due to Melrose '94 and Vasy '13-'19 (or assume  $g$  is a compactly supported perturbation of static metric) + **propagation estimates Hörmander '71**

## Positive commutator estimates

*Toy model:*  $P = P^*$  bounded, and  $\exists$  bounded  $A$  and  $D$  s.t.:

$$[P, iA] \geq (\mathbf{1} + D^2)^s. \quad (*)$$

Undo the commutator:

$$\begin{aligned} \frac{1}{2} \langle [P, iA]u, u \rangle &= \frac{\langle APu, u \rangle - \langle PAu, u \rangle}{2i} \\ &= \frac{\langle Pu, Au \rangle - \langle Au, Pu \rangle}{2i} \leq |\langle Pu, Au \rangle|, \end{aligned}$$

By Cauchy–Schwarz,

$$|\langle Pu, Au \rangle| \leq C \|(\mathbf{1} + D^2)^{-s/2} Pu\| \|(\mathbf{1} + D^2)^{s/2} u\| =: C \|Pu\|_{-s} \|u\|_s.$$

In combination with (\*):

$$\|u\|_s^2 \leq C \|Pu\|_{-s} \|u\|_s,$$

hence invertibility statement  $\|u\|_s \leq C \|Pu\|_{-s}$ .

# Positive commutator estimates

The existence of suitable  $A$  s.t.

$$[P, iA] \geq (\mathbf{1} + D^2)^s.$$

is **extremely rare**. But we can expect to prove it “somewhere in phase space”.

- ▶ If  $P \in \Psi^s(M)$  and  $A \in \Psi^\ell(M)$  then  $[P, iA] \in \Psi^{s+\ell-1}(M)$  and

$$\sigma_{\text{pr}}([P, iA]) = \{p, a\} \bmod S^{s+\ell-2}(M).$$

The flow of  $\{p, \cdot\}$  in  $\{p = 0\}$  is the classical Hamilton flow, or **bicharacteristic flow** (note that in  $\{p \neq 0\}$  elliptic theory applies).


- ▶ non-compact settings require weighted Sobolev spaces: extra weight  $(\mathbf{1} + |x|^2)^\ell$  ( $\Psi_{\text{sc}}^{m,\ell}(M)$  calculus)
- ▶ non-selfadjointness can be serious trouble (if we know nothing of  $P - P^*$ ), or valuable help (for instance  $P - i\varepsilon$  with  $\varepsilon > 0$ )

## Lorentzian scattering spaces

*Example:* Minkowski metric  $g_0 = dx_0^2 - (dx_1^2 + \dots + dx_{n-1}^2)$  on  $\mathbb{R}^n$  extends to **radial compactification**  $\overline{\mathbb{R}^n}$  defined using boundary-defining function  $\rho = (x_0^2 + x_1^2 + \dots + x_n^2)^{-\frac{1}{2}}$ . Regularity w.r.t.  $\rho^2 \partial_\rho = -\partial_r$

*Definition:* **Lorentzian sc-metrics** are  $C^\infty$  sections of  ${}^{\text{sc}}T^*M \otimes_s {}^{\text{sc}}T^*M$ , where  ${}^{\text{sc}}T^*M$  generated by  $\rho^{-2}d\rho, \rho^{-1}dy_1, \dots, \rho^{-1}dy_{n-1}$ .

Null geodesics lift to **null bicharacteristics** on  ${}^{\text{sc}}T^*M$  (rescaled and extended at  $\partial M$  appropriately)




*Definition:*

$(M, g)$  **non-trapping Lorentzian sc-space** if there are sinks/sources  $L_\pm$  above  $\partial M$ , and null bicharacteristics flow from and to  $L_-$  and  $L_+$ .

*Includes small perturbations of Minkowski space and asymptotically Minkowski spaces.*

## From null bicharacteristic flow to global estimates



dynamics of null bicharacteristics in  $\overline{scT^*M}$



classical quantities increasing along flow



pos. commutator estimates in  $\Psi_{sc}^{m,\ell}$ -calculus



1. Deduce **Fredholm property** and **invertibility** of  $P - z$
2. Deduce **singularities** of  $(P - z)^{-1}(x, x')$

# Dirac operators

The **Lorentzian Dirac operator**  $\mathbb{D}$  satisfies  $\mathbb{D}^2 = \square_g + \text{l.o.t.}$  in vector bundle sense. It is formally self-adjoint w.r.t. the canonical **indefinite** inner product, but (in general) not for an honest scalar product. However, on Lorentzian scattering spaces,  $P := \mathbb{D}^2$  satisfies

$$P^* - P \in \Psi_{\text{sc}}^{1, -1-\delta}(M)$$

for instance for the scalar product  $\langle \cdot, \gamma(n) \cdot \rangle_{L^2(M; SM)}$  used in quantization

✂ **work in progress** (with N.V. Dang & A. Vasy):  $P = \mathbb{D}^2$  on non-trapping Lorentzian scattering space  $(M, g)$  as closed operator.

## Conjecture

$\mathbb{D}^2$  is a closed operator, and:

$$\text{sp}(\mathbb{D}^2) \subset \mathbb{R} \cup \{\text{some isolated poles in } |\text{Im } z| \leq R\}$$

This uses stronger resolvent estimates using a resolved  $\Psi_{\text{sc}}^{m, \ell}$ -calculus obtained from blowing up the corner of  ${}^{\text{sc}}\overline{T^*M}$ .

# Dirac operators

The **Lorentzian Dirac operator**  $\mathcal{D}$  satisfies  $\mathcal{D}^2 = \square_g + \text{l.o.t.}$  in vector bundle sense. It is formally self-adjoint w.r.t. the canonical **indefinite** inner product, but (in general) not for an honest scalar product. However, on Lorentzian scattering spaces,  $P := \mathcal{D}^2$  satisfies

$$P^* - P \in \Psi_{\text{sc}}^{1, -1-\delta}(M)$$

for instance for the scalar product  $\langle \cdot, \gamma(n)\cdot \rangle_{L^2(M; SM)}$  used in quantization

✂ **work in progress** (with N.V. Dang & A. Vasy):  $P = \mathcal{D}^2$  on non-trapping Lorentzian scattering space  $(M, g)$  as closed operator.

## Conjecture

$\mathcal{D}^2$  is a closed operator, and:

$$\text{sp}(\mathcal{D}^2) \subset \mathbb{R} \cup \{\text{some isolated poles in } |\text{Im } z| \leq R\}$$

The techniques give a fully microlocal implementation of subelliptic estimate of Taira '21:


$$u \in H_{\text{sc}}^{m+\frac{1}{2}, \ell-\frac{1}{2}}(M), (P - z)u \in H_{\text{sc}}^{m, \ell}(M) \Rightarrow u \in H_{\text{sc}}^{m, \ell}(M).$$

# Dirac operators

The Lorentzian Dirac operator  $\mathcal{D}$  satisfies  $\mathcal{D}^2 = \square_g + \text{l.o.t.}$  in vector bundle sense. It is formally self-adjoint w.r.t. the canonical indefinite inner product, but (in general) not for an honest scalar product. However, on Lorentzian scattering spaces,  $P := \mathcal{D}^2$  satisfies

$$P^* - P \in \Psi_{\text{sc}}^{1, -1-\delta}(M)$$

for instance for the scalar product  $\langle \cdot, \gamma(n) \cdot \rangle_{L^2(M; SM)}$  used in quantization

 work in progress (with N.V. Dang & A. Vasy):  $P = \mathcal{D}^2$  on non-trapping Lorentzian scattering space  $(M, g)$  as closed operator.

## Conjecture

$\mathcal{D}^2$  is a closed operator, and:

$$\text{sp}(\mathcal{D}^2) \subset \mathbb{R} \cup \{\text{some isolated poles in } |\text{Im } z| \leq R\}$$

*Remark:* No role played by indefinite  $\langle \cdot, \cdot \rangle_{L^2(M; SM)}$



## **IV. Summary**

## To sum up...

We have shown relationship of Lorentzian spectral zeta function density  $\zeta_{g,\varepsilon}$  with space-time geometry.

$\Rightarrow$  (Lorentzian!) Gravity can be derived from a spectral action.

- 
- ▶ We also get the theorem for ultra-static spacetimes and compactly supported perturbations. One can conjecture extensions to asymptotically static spacetimes (and beyond, especially if weakening essential self-adjointness).
  - ▶ We show that the poles are a generalized Wodzicki residue
  - ▶ Relationships with QFT on curved spacetimes and renormalization
  - ▶  $(P - z)^{-1}$  contains information about null geodesics and causality
  - ② Does this fit into a spectral triple formalism? Non-commutative examples? Interpretation of spectrum?

*Thank you for your attention!*