

# HIDDEN GEOMETRY OF THE STANDARD MODEL

Andrzej Sitarz

(based on joint works with A. Bochniak & L. Dąbrowski)

Institute of Physics  
Jagiellonian University  
Kraków




Institute of Mathematics,  
Polish Academy of Sciences  
Warsaw



London, December 6th, 2018

# PHYSICS AND GEOMETRY



# SPIN GEOMETRY

## GEOMETRY OF FUNDAMENTAL PARTICLES

## GEOMETRY OF FUNDAMENTAL PARTICLES

- ① all fundamental fermions are described by spinor fields over a 4-dimensional Lorentzian manifold

## GEOMETRY OF FUNDAMENTAL PARTICLES

- 1 all fundamental fermions are described by spinor fields over a 4-dimensional Lorentzian manifold
- 2 spin geometry is seen through the existence of the Dirac operator constructed from torsion-free spin connection

## GEOMETRY OF FUNDAMENTAL PARTICLES

- 1 all fundamental fermions are described by spinor fields over a 4-dimensional Lorentzian manifold
- 2 spin geometry is seen through the existence of the Dirac operator constructed from torsion-free spin connection
- 3 spinors are sections of a complex vector bundle with a representation of the Clifford algebra

## GEOMETRY OF FUNDAMENTAL PARTICLES

- 1 all fundamental fermions are described by spinor fields over a 4-dimensional Lorentzian manifold
- 2 spin geometry is seen through the existence of the Dirac operator constructed from torsion-free spin connection
- 3 spinors are sections of a complex vector bundle with a representation of the Clifford algebra
- 4 the structure of the spinor bundle and the Dirac operator encodes the chirality and Lorentzian structure

## GEOMETRY OF FUNDAMENTAL PARTICLES

- 1 all fundamental fermions are described by spinor fields over a 4-dimensional Lorentzian manifold
- 2 spin geometry is seen through the existence of the Dirac operator constructed from torsion-free spin connection
- 3 spinors are sections of a complex vector bundle with a representation of the Clifford algebra
- 4 the structure of the spinor bundle and the Dirac operator encodes the chirality and Lorentzian structure
- 5 the spinor bundle of fermions is an irreducible representation of the chiral Clifford algebra (and that can be encoded in saying that the bundle establishes the Morita equivalence between functions and Clifford algebra).

# SPECTRAL TRIPLES

## DEFINITION: A GEOMETRY (ALGEBRAIC WAY)

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint operator  $D$ , satisfying several conditions.

- 1 make sure that  $D$  is a first-order operator

# SPECTRAL TRIPLES

## DEFINITION: A GEOMETRY (ALGEBRAIC WAY)

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint operator  $D$ , satisfying several conditions.

- 1 make sure that  $D$  is a first-order operator
- 2 choose  $D$  to be nondegenerate,

# SPECTRAL TRIPLES

## DEFINITION: A GEOMETRY (ALGEBRAIC WAY)

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint operator  $D$ , satisfying several conditions.

- 1 make sure that  $D$  is a first-order operator
- 2 choose  $D$  to be nondegenerate,
- 3 guarantee that  $D$  is a differential operator
- 4 try to define the dimension & ...+ make sure that you can compute something

# SPECTRAL TRIPLES

## DEFINITION: A GEOMETRY (ALGEBRAIC WAY)

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint operator  $D$ , satisfying several conditions.

- 1 make sure that  $D$  is a first-order operator
- 2 choose  $D$  to be nondegenerate,
- 3 guarantee that  $D$  is a differential operator
- 4 try to define the dimension & ...+ make sure that you can compute something

## THEOREM [CONNES]

If you have a spin manifold then the Dirac operator acting on the sections of the spinor bundle gives you geometry in the above sense.

# SPECTRAL TRIPLES

## DEFINITION: A GEOMETRY (ALGEBRAIC WAY)

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint operator  $D$ , satisfying several conditions.

- 1 make sure that  $D$  is a first-order operator
- 2 choose  $D$  to be nondegenerate,
- 3 guarantee that  $D$  is a differential operator
- 4 try to define the dimension & ...+ make sure that you can compute something

## THEOREM [CONNES]

If you have a spin manifold then the Dirac operator acting on the sections of the spinor bundle gives you geometry in the above sense.

## REMARK

Another natural triple is  $(C^\infty(M), L^2(\Omega(M)), d + d^*)$ .

# THE STRUCTURE OF THE STANDARD MODEL

## TESTING THE EUCLIDEAN SM

# THE STRUCTURE OF THE STANDARD MODEL

## TESTING THE EUCLIDEAN SM

- ① Each fermion is described by a spectral triple (spinor + Dirac)

# THE STRUCTURE OF THE STANDARD MODEL

## TESTING THE EUCLIDEAN SM

- 1 Each fermion is described by a spectral triple (spinor + Dirac)
- 2 the collection of all particles with the interactions corresponds to a finite spectral triple

# THE STRUCTURE OF THE STANDARD MODEL

## TESTING THE EUCLIDEAN SM

- 1 Each fermion is described by a spectral triple (spinor + Dirac)
- 2 the collection of all particles with the interactions corresponds to a finite spectral triple
- 3 the full geometry of SM is (almost) a product geometry

# THE STRUCTURE OF THE STANDARD MODEL

## TESTING THE EUCLIDEAN SM

- 1 Each fermion is described by a spectral triple (spinor + Dirac)
- 2 the collection of all particles with the interactions corresponds to a finite spectral triple
- 3 the full geometry of SM is (almost) a product geometry
- 4 the fluctuations of the Dirac operator are gauge fields

# THE STRUCTURE OF THE STANDARD MODEL

## TESTING THE EUCLIDEAN SM

- 1 Each fermion is described by a spectral triple (spinor + Dirac)
- 2 the collection of all particles with the interactions corresponds to a finite spectral triple
- 3 the full geometry of SM is (almost) a product geometry
- 4 the fluctuations of the Dirac operator are gauge fields

## FINITE SPECTRAL TRIPLE OF THE SM

The “almost commutative” geometry is described by the Hilbert space

$$H_F = \mathbb{C}^{96} =: H_f \otimes \mathbb{C}^3,$$

$$\begin{bmatrix} \nu_R & u_R^1 & u_R^2 & u_R^3 \\ e_R & d_R^1 & d_R^2 & d_R^3 \\ \nu_L & u_L^1 & u_L^2 & u_L^3 \\ e_L & d_L^1 & d_L^2 & d_L^3 \end{bmatrix} \quad \begin{bmatrix} \bar{\nu}_R & \bar{e}_R & \bar{\nu}_L & \bar{e}_L \\ \bar{u}_R^1 & \bar{d}_R^1 & \bar{u}_L^1 & \bar{d}_L^1 \\ \bar{u}_R^2 & \bar{d}_R^2 & \bar{u}_L^2 & \bar{d}_L^2 \\ \bar{u}_R^3 & \bar{d}_R^3 & \bar{u}_L^3 & \bar{d}_L^3 \end{bmatrix}$$

# THE SPECTRAL TRIPLE OF THE SM

## THE ALGEBRA FOR $g$ GENERATIONS

$$A_F = (M_1 \oplus M_1 \oplus M_2)^{(4g)} \oplus (M_1 \oplus M_3)^{(4g)},$$

with first  $M_1$  identified with each other:

$$(z \oplus w \oplus h)^{(4g)} \oplus (z \oplus m)^{(4g)}.$$

# THE SPECTRAL TRIPLE OF THE SM

## THE ALGEBRA FOR $g$ GENERATIONS

$$A_F = (M_1 \oplus M_1 \oplus M_2)^{(4g)} \oplus (M_1 \oplus M_3)^{(4g)},$$

with first  $M_1$  identified with each other:

$$(z \oplus w \oplus h)^{(4g)} \oplus (z \oplus m)^{(4g)}.$$

## THE DIRAC OPERATOR

$$\tilde{D}_F = (D_l \oplus D_q^{(3)}) \oplus 0^{(16g)},$$

# THE DIRAC OPERATOR AND HODGE DUALITY

## THE DIRAC OPERATOR

$$D_l = \left( \begin{array}{cc|cc} 0 & 0 & \gamma_\nu & 0 \\ 0 & 0 & 0 & \gamma_e \\ \hline \gamma_\nu^* & 0 & 0 & 0 \\ 0 & \gamma_e^* & 0 & 0 \end{array} \right), \quad D_q = \left( \begin{array}{cc|cc} 0 & 0 & \gamma_u & 0 \\ 0 & 0 & 0 & \gamma_d \\ \hline \gamma_u^* & 0 & 0 & 0 \\ 0 & \gamma_d^* & 0 & 0 \end{array} \right),$$

where where  $D_l, D_q \in M_{4g}$  are positive mass matrices for leptons and quarks, respectively.

# THE DIRAC OPERATOR AND HODGE DUALITY

## THE DIRAC OPERATOR

$$D_l = \left( \begin{array}{cc|cc} 0 & 0 & \gamma_\nu & 0 \\ 0 & 0 & 0 & \gamma_e \\ \hline \gamma_\nu^* & 0 & 0 & 0 \\ 0 & \gamma_e^* & 0 & 0 \end{array} \right), \quad D_q = \left( \begin{array}{cc|cc} 0 & 0 & \gamma_u & 0 \\ 0 & 0 & 0 & \gamma_d \\ \hline \gamma_u^* & 0 & 0 & 0 \\ 0 & \gamma_d^* & 0 & 0 \end{array} \right),$$

where where  $D_l, D_q \in M_{4g}$  are positive mass matrices for leptons and quarks, respectively.

## QUESTION

What is the algebra generated by  $\tilde{D}_F$  and  $A_F$  and what is its commutant? If the commutant of  $Cl(A_F, D_F)$  is isomorphic to  $A_F$  then SM corresponds to fundamental spinors. *BUT...*

# THE HODGE DUALITY

## SM AS GEOMETRY

If the algebra generated by  $A_F, \tilde{D}_F$  is

$$M_{4g} \oplus M_{4g}^3 \oplus M_1^{4g} \oplus M_3^{4g}$$

then the geometry is self-dual (commutant of the Clifford algebra is the Clifford algebra itself)

# THE HODGE DUALITY

## SM AS GEOMETRY

If the algebra generated by  $A_F, \tilde{D}_F$  is

$$M_{4g} \oplus M_{4g}^3 \oplus M_1^{4g} \oplus M_3^{4g}$$

then the geometry is self-dual (commutant of the Clifford algebra is the Clifford algebra itself)

## DOES IT HAPPEN IN REALITY?

Yes, if there is nontrivial mixing of the neutrino flavors and nontrivial mixing of the quarks and all fermion (bare) masses are nonzero and different from each other then the SM satisfies the Hodge duality.

# THE DETAILS

## ONE GENERATION

In this case the various  $\Upsilon$ 's are just complex numbers and so the  $D_l$  and  $D_q$  matrices are just  $4 \times 4$  complex matrices acting on the spaces of leptons and quarks, where  $\Upsilon_e$  is the electron mass and  $\Upsilon_\nu$  is the neutrino mass (and similarly for  $D_q$ ).

Since  $\mathbb{C}l_D(A)$  contains  $M_1^{(4)} \oplus M_3^{(4)}$ , the commutant  $\mathbb{C}l_D(A)'$  of  $\mathbb{C}l_D(A)$  must contain  $M_4 \oplus M_4^{(3)}$ , and, if the Hodge duality is satisfied, so must  $\mathbb{C}l_D(A)$ . However,  $\mathbb{C}l_D(A)$  contains  $M_1^{(4)} \oplus M_3^{(4)}$  and two algebras generated respectively by  $M_1 \oplus M_1 \oplus M_2$  and  $D_l$ , and  $(M_1 \oplus M_1 \oplus M_2)^{(3)}$  and  $D_q^{(3)}$ . Thus the only possibility that the Hodge condition holds is when these two algebras are  $M_4$  and  $M_4^{(3)}$ , respectively. This happens only when independently all  $z \oplus w \oplus h \in M_1 \oplus M_1 \oplus M_2$  and  $D_l$ , as well as  $z \oplus w \oplus h$  and  $D_q$ , each generate  $M_4$ . It is easy to notice that sufficient and necessary condition for this is that all four masses  $\Upsilon$ 's are different from zero.

# THREE GENERATIONS

## LEPTONS

We start with leptons and check whether the algebra generated by  $A_l, D_l$  is a full matrix algebra. A general matrix that commutes with  $A_l$  has a form  $P_1 \oplus P_2 \oplus \tilde{P}_3$ , where  $P_1, P_2 \in M_g$  and  $\tilde{P}_3 = 1 \otimes P_3 \in M_2 \otimes M_g$ . If it commutes with  $D_l$  then:

$$\begin{aligned} P_1 \Upsilon_\nu &= \Upsilon_\nu P_3, & P_2 \Upsilon_e &= \Upsilon_e P_3, \\ P_3 \Upsilon_\nu^* &= \Upsilon_\nu^* P_1, & P_3 \Upsilon_e^* &= \Upsilon_e^* P_1. \end{aligned}$$

From these equations we immediately infer that  $P_1$  and  $P_3$  must commute with  $\Upsilon_\nu \Upsilon_\nu^*$  (note that since both  $\Upsilon$  matrices are unitarily similar to diagonal matrix then they are normal) whereas  $P_2$  and  $P_3$  must commute with  $\Upsilon_e \Upsilon_e^*$ .

# THREE GENERATIONS

## ...AND QUARKS

Therefore, only if both  $\Upsilon_\nu$  and  $\Upsilon_e$  are invertible and the pair  $\Upsilon_\nu \Upsilon_\nu^*$  and  $\Upsilon_e \Upsilon_e^*$  generate the full matrix algebra  $M_g$  it follows consequently that  $P_1$  and  $P_2$  must be equal to  $P_3$ , and be proportional to the identity. Similar arguments will also hold for the quarks: it suffices (and is necessary) that the two matrices  $\Upsilon_u \Upsilon_u^*$  and  $\Upsilon_d \Upsilon_d^*$  generate the full matrix algebra  $M_g$  and that they are invertible to assure that the algebra generated by  $A_q, D_q$  is a full matrix algebra.

# THREE GENERATIONS

## MASSES

When two hermitian matrices,  $A, B$  in  $M_3(\mathbb{C})$  generate a full matrix algebra ?

Burnside (1905): they do not share a common eigenvector (the theorem states that there is no common invariant subspace but since the matrices are hermitian if there exists an invariant subspace its complement is also invariant and hence there would necessarily exist an invariant subspace of dimension 1).

If, we assume that all eigenvalues of  $A$  are different from each other then we only need to check the matrix elements of  $U$  in the chosen basis, in which  $A$  is diagonal. If no matrix element of  $U$  is of modulus 1 (while at the same time other matrix elements in the same row and in the same column are 0) then  $U$  does not map one of the basis vectors to another one. Equivalently, one can reformulate the condition in the following way: no permutation of the basis leads to the block diagonal matrix of  $U$  with rank of the largest block strictly less than 3.

## THE PHYSICAL PARAMETERS

$$U = \begin{bmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{bmatrix},$$

where  $s_{ij} = \sin \theta_{ij}$  and  $c_{ij} = \cos \theta_{ij}$ . The following Jarlskog invariant:

$$J_{\nu CP}^{max} = \frac{1}{8} \cos(\theta_{13}) \sin(2\theta_{13}) \sin(2\theta_{23}) \sin(2\theta_{12}),$$

provides for the neutrino mixing range (with  $1\sigma$  error):

$$J_{\nu CP}^{max} = 0.0329 \pm 0.0007,$$

that proves indeed that not only all the angles are non-vanishing but all neutrino masses are different from each other.

## THE PHYSICAL PARAMETERS: QUARKS

Here the usual convention in physics is different with "up" sector diagonal and "down" sector mixed. The bare up and down quark masses are different from each other within the errors so the only thing to check is the mixing matrix  $U$ . Using again the same type of parametrization of the matrix by three angles and the phases, we can just look at the experimental value of the Jarlskog invariant  $J_{qCP}^{max}$ , measured with  $1\sigma$ :

$$J_{qCP}^{max} = \left( 3.04 \begin{matrix} +0.21 \\ -0.20 \end{matrix} \right) 10^{-5},$$

which is sufficient to ascertain that all angles are indeed nonzero and that implies the partial condition to Hodge duality.

Note that unlike in the leptonic case the angles are very small, which means that the matrix  $U$  is very close to the diagonal unit matrix. Nevertheless within the experimental errors we see that the quark mixing is also maximal and the Clifford algebra generated in the quark sector is the full matrix algebra  $M_{4g}$  as well.

# THE LORENTZIAN SPECTRAL TRIPLES

## THE DEFINITION

A real pseudo-Riemannian spectral triple of signature  $(p, q)$  is a system  $(\mathcal{A}, \pi, \mathcal{H}, D, J, \gamma, \beta)$  where  $\mathcal{A}$  is an involutive unital algebra,  $\pi$  its faithful  $*$ -representation on an Hilbert space  $\mathcal{H}$ . First, for even  $p+q$  there exists a  $\mathbb{Z}_2$ -grading  $\gamma^\dagger = \gamma, \gamma^2 = 1$  commuting with the representation of  $\mathcal{A}$ ,  $J$  is an antilinear isometry and for all  $a, b \in \mathcal{A}$  we have  $[J\pi(a^*)J^{-1}, \pi(b)] = 0$ . Furthermore, there exists an additional grading  $\beta = \beta^\dagger, \beta^2 = 1$  also commuting with the representation of  $\mathcal{A}$ , which defines the Krein structure on the Hilbert space. The latter is an indefinite bilinear form defined as  $(\phi, \psi)_\beta = (\phi, \beta\psi)$ , where  $(\cdot, \cdot)$  is the usual positive definite scalar product on the Hilbert space. As a last requirement, we postulate the existence of a (possibly unbounded) densely defined operator  $D$ , which is  $\beta$ -self-adjoint, i.e.  $D^\dagger = (-1)^p \beta D \beta$  and such that  $[D, \pi(a)]$  is bounded for every  $a \in \mathcal{A}$ , is odd with respect to  $\gamma$ -grading:  $D\gamma = -\gamma D$ .

# FINITE LORENTZIAN TRIPLES

## THE EUCLIDEAN PART OF FST

Define  $D_+ = \frac{1}{2}(D + D^\dagger)$  and  $D_- = \frac{i}{2}(D - D^\dagger)$ . Both  $D_\pm$  are by definition self-adjoint and we obtain a pair of Riemannian real spectral triples,  $(\mathcal{A}, \pi, \mathcal{H}, D_\pm, J, \gamma)$ .

# FINITE LORENTZIAN TRIPLES

## THE EUCLIDEAN PART OF FST

Define  $D_+ = \frac{1}{2}(D + D^\dagger)$  and  $D_- = \frac{i}{2}(D - D^\dagger)$ . Both  $D_\pm$  are by definition self-adjoint and we obtain a pair of Riemannian real spectral triples,  $(\mathcal{A}, \pi, \mathcal{H}, D_\pm, J, \gamma)$ .

## MAIN FEATURES OF FST

# FINITE LORENTZIAN TRIPLES

## THE EUCLIDEAN PART OF FST

Define  $D_+ = \frac{1}{2}(D + D^\dagger)$  and  $D_- = \frac{i}{2}(D - D^\dagger)$ . Both  $D_\pm$  are by definition self-adjoint and we obtain a pair of Riemannian real spectral triples,  $(\mathcal{A}, \pi, \mathcal{H}, D_\pm, J, \gamma)$ .

## MAIN FEATURES OF FST

- 1 The classification of Dirac operator is almost the same as in the Riemannian case.

# FINITE LORENTZIAN TRIPLES

## THE EUCLIDEAN PART OF FST

Define  $D_+ = \frac{1}{2}(D + D^\dagger)$  and  $D_- = \frac{i}{2}(D - D^\dagger)$ . Both  $D_\pm$  are by definition self-adjoint and we obtain a pair of Riemannian real spectral triples,  $(\mathcal{A}, \pi, \mathcal{H}, D_\pm, J, \gamma)$ .

## MAIN FEATURES OF FST

- 1 The classification of Dirac operator is almost the same as in the Riemannian case.
- 2 The existence of  $\beta$  and the time-orientability puts further restrictions.

# FINITE LORENTZIAN TRIPLES

## THE EUCLIDEAN PART OF FST

Define  $D_+ = \frac{1}{2}(D + D^\dagger)$  and  $D_- = \frac{i}{2}(D - D^\dagger)$ . Both  $D_\pm$  are by definition self-adjoint and we obtain a pair of Riemannian real spectral triples,  $(\mathcal{A}, \pi, \mathcal{H}, D_\pm, J, \gamma)$ .

## MAIN FEATURES OF FST

- 1 The classification of Dirac operator is almost the same as in the Riemannian case.
- 2 The existence of  $\beta$  and the time-orientability puts further restrictions.
- 3 If for an even spectral triple grading is scalar on at least one subspace  $H_{ij}$  then there is no pseudo-Riemannian triple with  $p$  odd.

# FINITE LORENTZIAN TRIPLES

## THE EUCLIDEAN PART OF FST

Define  $D_+ = \frac{1}{2}(D + D^\dagger)$  and  $D_- = \frac{i}{2}(D - D^\dagger)$ . Both  $D_\pm$  are by definition self-adjoint and we obtain a pair of Riemannian real spectral triples,  $(\mathcal{A}, \pi, \mathcal{H}, D_\pm, J, \gamma)$ .

## MAIN FEATURES OF FST

- 1 The classification of Dirac operator is almost the same as in the Riemannian case.
- 2 The existence of  $\beta$  and the time-orientability puts further restrictions.
- 3 If for an even spectral triple grading is scalar on at least one subspace  $H_{ij}$  then there is no pseudo-Riemannian triple with  $p$  odd.
- 4 The Euclidean part of ST is a finite spectral triple **BUT** with an additional symmetry  $\beta$  of the Dirac operator.

# THE LEPTON NUMBER SYMMETRY FROM LORENTZIAN STRUCTURE

## THE PSEUDO-RIEMANNIAN STRUCTURE

The finite spectral triple of the Standard Model is consistent with it being the Euclidean part of the pseudo-Riemannian triple with  $\beta$  being:

$$\beta = \pi_F(1, 1, -1)J_F\pi_F(1, 1, -1)J_F^{-1},$$

where  $\lambda, h, m \in M_1 \oplus M_2 \oplus M_3$ .

# THE LEPTON NUMBER SYMMETRY FROM LORENTZIAN STRUCTURE

## THE PSEUDO-RIEMANNIAN STRUCTURE

The finite spectral triple of the Standard Model is consistent with it being the Euclidean part of the pseudo-Riemannian triple with  $\beta$  being:

$$\beta = \pi_F(1, 1, -1)J_F\pi_F(1, 1, -1)J_F^{-1},$$

where  $\lambda, h, m \in M_1 \oplus M_2 \oplus M_3$ .

## IS THIS UNIQUE ?

Yes, it is the only possible **0**-cycle ( $\beta$ ) for a real spectral triple over the Standard Model that can be interpreted shadow of a pseudo-Riemannian structure, which additionally allows Hodge duality is the one with  $\beta = \pi(1, 1, -1)J\pi(1, 1, -1)J^{-1}$ .

This results in the symmetry that physically can be interpreted as **lepton number conservation**.

# CONCLUSIONS

- 1 The finite structure (particle contents) of the Standard Model has some interesting geometric features

# CONCLUSIONS

- 1 The finite structure (particle contents) of the Standard Model has some interesting geometric features
- 2 Particles correspond rather to **forms** than **spinors**

# CONCLUSIONS

- ① The finite structure (particle contents) of the Standard Model has some interesting geometric features
- ② Particles correspond rather to **forms** than **spinors**
- ③ Hodge duality is satisfied for generations **but** requires mixing of flavors both in the leptonic as well as quark sector.

# CONCLUSIONS

- 1 The finite structure (particle contents) of the Standard Model has some interesting geometric features
- 2 Particles correspond rather to **forms** than **spinors**
- 3 Hodge duality is satisfied for generations **but** requires mixing of flavors both in the leptonic as well as quark sector.
- 4 The model looks like a Euclidean restriction of the pseudo-Riemannian one and the resulting symmetry preserves the lepton number **and** prevents breaking of the  **$SU(3)$**  symmetry

# CONCLUSIONS

- 1 The finite structure (particle contents) of the Standard Model has some interesting geometric features
- 2 Particles correspond rather to **forms** than **spinors**
- 3 Hodge duality is satisfied for generations **but** requires mixing of flavors both in the leptonic as well as quark sector.
- 4 The model looks like a Euclidean restriction of the pseudo-Riemannian one and the resulting symmetry preserves the lepton number **and** prevents breaking of the  **$SU(3)$**  symmetry

## LITERATURE

A. Bochniak, A.Sitarz, Finite Pseudo-Riemannian Spectral Triples and The Standard Model Phys. Rev. D 97, 115029 (2018)

L.Dąbrowski, A.Sitarz, Fermion masses, mass-mixing and the almost commutative geometry of the Standard Model, arXiv:1806.07282

# CONCLUSIONS

- 1 The finite structure (particle contents) of the Standard Model has some interesting geometric features
- 2 Particles correspond rather to **forms** than **spinors**
- 3 Hodge duality is satisfied for generations **but** requires mixing of flavors both in the leptonic as well as quark sector.
- 4 The model looks like a Euclidean restriction of the pseudo-Riemannian one and the resulting symmetry preserves the lepton number **and** prevents breaking of the  **$SU(3)$**  symmetry

## LITERATURE

A. Bochniak, A.Sitarz, Finite Pseudo-Riemannian Spectral Triples and The Standard Model Phys. Rev. D 97, 115029 (2018)

L.Dąbrowski, A.Sitarz, Fermion masses, mass-mixing and the almost commutative geometry of the Standard Model, arXiv:1806.07282

## THANK YOU