

# POINTLESS GEOMETRY

Andrzej Sitarz



Kraków, 10.03.2016

# GEOMETRY AND PHYSICS

## A MARRIAGE MADE IN HEAVEN

# GEOMETRY AND PHYSICS

## A MARRIAGE MADE IN HEAVEN

- early 20th century: gravitation and electromagnetism formulated as field theories in four-dimensional space-time,

# GEOMETRY AND PHYSICS

## A MARRIAGE MADE IN HEAVEN

- early 20th century: gravitation and electromagnetism formulated as field theories in four-dimensional space-time,
- early 20th century: quantum physics formulated using the analysis of linear operators and spectral theory

# GEOMETRY AND PHYSICS

## A MARRIAGE MADE IN HEAVEN

- early 20th century: gravitation and electromagnetism formulated as field theories in four-dimensional space-time,
- early 20th century: quantum physics formulated using the analysis of linear operators and spectral theory
- 1955: Yang-Mills theory - geometry (vector bundles, connections, curvatures, covariant derivatives and spinors)

# GEOMETRY AND PHYSICS

## A MARRIAGE MADE IN HEAVEN

- early 20th century: gravitation and electromagnetism formulated as field theories in four-dimensional space-time,
- early 20th century: quantum physics formulated using the analysis of linear operators and spectral theory
- 1955: Yang-Mills theory - geometry (vector bundles, connections, curvatures, covariant derivatives and spinors)

## ATIYAH, DIJKGRAAF, HITCHIN

Recent ideas from quantum gravity and string theory challenge the fundamental concepts of geometry at an even deeper level. Physical intuition tells us that the traditional pseudo-Riemannian geometry of space-time cannot be a definite description of physical reality.

# GEOMETRY AND PHYSICS

## THE FUTURE:

The idea is that the classical laws of gravity (and classical field theories) only appear in some limit, very similar to the emergence of the macroscopic laws of thermodynamics out of the microscopic description of statistical mechanics. The definite mathematical formulation of such a concept of "quantum geometry" is still far away.

# GEOMETRY AND PHYSICS

## THE FUTURE:

The idea is that the classical laws of gravity (and classical field theories) only appear in some limit, very similar to the emergence of the macroscopic laws of thermodynamics out of the microscopic description of statistical mechanics. The definite mathematical formulation of such a concept of "quantum geometry" is still far away.

## THE QUANTUM GEOMETRY

There are numerous ways and approaches to quantum geometry . Even the phrase noncommutative geometry has several interpretations.

# GEOMETRY AND PHYSICS

## THE FUTURE:

The idea is that the classical laws of gravity (and classical field theories) only appear in some limit, very similar to the emergence of the macroscopic laws of thermodynamics out of the microscopic description of statistical mechanics. The definite mathematical formulation of such a concept of "quantum geometry" is still far away.

## THE QUANTUM GEOMETRY

There are numerous ways and approaches to quantum geometry . Even the phrase noncommutative geometry has several interpretations.

## THE QUESTION ?

Is there a way to describe geometry using the methods, which do not use the existence of points ?

# GEOMETRY WITHOUT POINTS

HOW DO WE SEE A POINT IN A SPACE?

# GEOMETRY WITHOUT POINTS

HOW DO WE SEE A POINT IN A SPACE?

We don't. We see and „measure“ **functions** on the space.

# GEOMETRY WITHOUT POINTS

## HOW DO WE SEE A POINT IN A SPACE?

We don't. We see and „measure“ **functions** on the space.

## EXAMPLE

If  $X$  is a (locally) compact Hausdorff space and  $C(X)$  is the algebra of continuous functions on  $X$ , then  $C(X)$  is a commutative (non) unital  $C^*$ -algebra.

# GEOMETRY WITHOUT POINTS

## HOW DO WE SEE A POINT IN A SPACE?

We don't. We see and „measure“ **functions** on the space.

## EXAMPLE

If  $X$  is a (locally) compact Hausdorff space and  $C(X)$  is the algebra of continuous functions on  $X$ , then  $C(X)$  is a commutative (non) unital  $C^*$ -algebra.

## WHAT IS A $C^*$ -ALGEBRA?

Let  $\mathcal{A}$  be an involutive Banach algebra (that is a complex normed algebra, which is complete as a topological space in the norm). If:

$$\|aa^*\| = \|a\|^2,$$

then  $\mathcal{A}$  is a  $C^*$ -algebra.

# GELFAND-NAIMARK THEOREMS

## THEOREM (GELFAND-NAIMARK (GNS))

*Every abstract  $C^*$ -algebra  $\mathcal{A}$  is isometrically  $*$ -isomorphic to a concrete  $C^*$  algebra of operators on a Hilbert space  $\mathcal{H}$ . If the algebra  $\mathcal{A}$  is separable then we can take  $\mathcal{H}$  to be separable.*

# GELFAND-NAIMARK THEOREMS

## THEOREM (GELFAND-NAIMARK (GNS))

*Every abstract  $C^*$ -algebra  $\mathcal{A}$  is isometrically  $*$ -isomorphic to a concrete  $C^*$  algebra of operators on a Hilbert space  $\mathcal{H}$ . If the algebra  $\mathcal{A}$  is separable then we can take  $\mathcal{H}$  to be separable.*

## THEOREM (GELFAND-NAIMARK (GN))

*If a  $C^*$  algebra is commutative then the Gelfand transform,  $\mu : \mathcal{A} \rightarrow C_0(X_{\mathcal{A}})$ , where  $X_{\mathcal{A}}$  is the Gelfand spectrum of  $\mathcal{A}$  (space of homomorphisms from  $\mathcal{A}$  to  $\mathbb{C}$ ):*

$$\mu(a)(\chi) = \chi(a), \quad a \in \mathcal{A}, \chi \in X_{\mathcal{A}},$$

*is an isometric  $*$ -isomorphism.*

# WHAT IS NONCOMMUTATIVE TOPOLOGY

## TOPOLOGY vs ALGEBRA

(locally compact) topological space  
homeomorphism  
continuous proper map  
compact space  
open (dense) subset  
compactification  
Stone-Čech compactification  
cartesian product

$C^*$ -algebra  
automorphism  
morphism  
unital  $C^*$ -algebra  
(essential) ideal  
unitization  
multiplier algebra  
tensor product

### IS THAT ENOUGH?

# WHAT IS NONCOMMUTATIVE TOPOLOGY

## TOPOLOGY vs ALGEBRA

(locally compact) topological space	$C^*$ -algebra
homeomorphism	automorphism
continuous proper map	morphism
compact space	unital $C^*$ -algebra
open (dense) subset	(essential) ideal
compactification	unitization
Stone-Čech compactification	multiplier algebra
cartesian product	tensor product

## IS THAT ENOUGH?

There is much more to **geometry** than topology

# WHAT IS NONCOMMUTATIVE TOPOLOGY

## TOPOLOGY vs ALGEBRA

(locally compact) topological space	$C^*$ -algebra
homeomorphism	automorphism
continuous proper map	morphism
compact space	unital $C^*$ -algebra
open (dense) subset	(essential) ideal
compactification	unitization
Stone-Čech compactification	multiplier algebra
cartesian product	tensor product

## IS THAT ENOUGH?

There is much more to **geometry** than topology

There are **groups**, **vector bundles**, **connections** and **metric**.

# THE FINAL DICTIONARY


## GEOMETRY

vector bundle  
differential forms  
differential forms  
de Rham cohomology  
vector fields  
group  
Lie algebra  
principal fibre bundle  
measurable functions  
infinitesimals  
metric  
spin<sup>c</sup> geometry  
spin geometry  
integrals


## ALGEBRA

finitely generated projective module  
differential forms  
Hochschild homology  
cyclic cohomology  
operators  
Hopf algebra  
Hopf algebra  
Hopf-Galois extension  
von Neumann algebra  
compact operators  
Dirac operator  
spectral triple  
real spectral triple  
exotic traces

# THE IMAGE OF NONCOMMUTATIVE GEOMETRY




# THE IMAGE OF NONCOMMUTATIVE GEOMETRY



IS THERE A GAIN FOR PHYSICS ?

# THE IMAGE OF NONCOMMUTATIVE GEOMETRY



IS THERE A GAIN FOR PHYSICS ?

**YES!** We have a much broader interpretation of what **geometry** is and we can look at what is observed from a completely different point of view.

# THE DIRAC OPERATOR

## FORGET THE USUAL CONSTRUCTION

Riemannian manifold (compact, closed) with a metric  $g$ . Find Clifford algebra bundle, identify the spinor bundle, then lift the metric connection (the Levi-Civita one if you want the **true** Dirac), lift it to the spinor bundle, compose with Clifford map – you get  $D$ . Then prove some theorems about  $D$ .

# THE DIRAC OPERATOR

## FORGET THE USUAL CONSTRUCTION

Riemannian manifold (compact, closed) with a metric  $g$ . Find Clifford algebra bundle, identify the spinor bundle, then lift the metric connection (the Levi-Civita one if you want the **true** Dirac), lift it to the spinor bundle, compose with Clifford map – you get  $D$ . Then prove some theorems about  $D$ .

## FOLLOW THE OPERATIONAL DEFINITION

Take an algebra represented on a Hilbert space and look for operators, which **behave** like Dirac operators: first order differential operator such that gives you the metric:

$$d(x, y) = \sup_{\| [D, f] \| \leq 1} |f(x) - f(y)|,$$

is unbounded, discrete spectrum and the eigenvalues have certain growth.

# GEOMETRY AND THE HILBERT SPACES.

## THE SIGNIFICANCE OF ATIYAH-SINGER

All **classical** geometry can be encoded in terms of operators on a separable Hilbert space.

# GEOMETRY AND THE HILBERT SPACES.

## THE SIGNIFICANCE OF ATIYAH-SINGER

All **classical** geometry can be encoded in terms of operators on a separable Hilbert space.

## HOW DO WE **RECONSTRUCT** GEOMETRY ?

# GEOMETRY AND THE HILBERT SPACES.

## THE SIGNIFICANCE OF ATIYAH-SINGER

All **classical** geometry can be encoded in terms of operators on a separable Hilbert space.

## HOW DO WE **RECONSTRUCT** GEOMETRY ?

- ① differential calculus:  $da = [D, a]$

# GEOMETRY AND THE HILBERT SPACES.

## THE SIGNIFICANCE OF ATIYAH-SINGER

All **classical** geometry can be encoded in terms of operators on a separable Hilbert space.

## HOW DO WE **RECONSTRUCT** GEOMETRY ?

- ① differential calculus:  $da = [D, a]$
- ② metric:  $d(x, y) = \sup_{\|[D, f]\| \leq 1} |f(x) - f(y)|$

# GEOMETRY AND THE HILBERT SPACES.

## THE SIGNIFICANCE OF ATIYAH-SINGER

All **classical** geometry can be encoded in terms of operators on a separable Hilbert space.

## HOW DO WE **RECONSTRUCT** GEOMETRY ?

- 1 differential calculus:  $da = [D, a]$
- 2 metric:  $d(x, y) = \sup_{\|[D, f]\| \leq 1} |f(x) - f(y)|$
- 3 additional connection (if spinors twisted by a vector bundle)

# GEOMETRY AND THE HILBERT SPACES.

## THE SIGNIFICANCE OF ATIYAH-SINGER

All **classical** geometry can be encoded in terms of operators on a separable Hilbert space.

## HOW DO WE **RECONSTRUCT** GEOMETRY ?

- 1 differential calculus:  $da = [D, a]$
- 2 metric:  $d(x, y) = \sup_{\| [D, f] \| \leq 1} |f(x) - f(y)|$
- 3 additional connection (if spinors twisted by a vector bundle)
- 4 dimension (growth of eigenvalues:  $N(\Lambda) \sim \Lambda^d$ ),

# GEOMETRY AND THE HILBERT SPACES.

## THE SIGNIFICANCE OF ATIYAH-SINGER

All **classical** geometry can be encoded in terms of operators on a separable Hilbert space.

## HOW DO WE **RECONSTRUCT** GEOMETRY ?

- 1 differential calculus:  $da = [D, a]$
- 2 metric:  $d(x, y) = \sup_{\| [D, f] \| \leq 1} |f(x) - f(y)|$
- 3 additional connection (if spinors twisted by a vector bundle)
- 4 dimension (growth of eigenvalues:  $N(\Lambda) \sim \Lambda^d$ ),
- 5 integral (exotic traces)

$$\text{Tr}(a) = \text{Res}_{z=d} \text{tr}(a|D|^{-z})$$

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

- We do not need **points** and manifolds.

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

- We do not need **points** and manifolds.
- We look for representations of **algebras** rather than **groups**.

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

- We do not need **points** and manifolds.
- We look for representations of **algebras** rather than **groups**.
- Symmetries are easily generalized in terms of **Hopf algebras**.

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

- We do not need **points** and manifolds.
- We look for representations of **algebras** rather than **groups**.
- Symmetries are easily generalized in terms of **Hopf algebras**.
- Metric and gauge fields are unified in an elegant way.

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

- We do not need **points** and manifolds.
- We look for representations of **algebras** rather than **groups**.
- Symmetries are easily generalized in terms of **Hopf algebras**.
- Metric and gauge fields are unified in an elegant way.
- The theory is not abstract (not **formal** manipulations).

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

- We do not need **points** and manifolds.
- We look for representations of **algebras** rather than **groups**.
- Symmetries are easily generalized in terms of **Hopf algebras**.
- Metric and gauge fields are unified in an elegant way.
- The theory is not abstract (not **formal** manipulations).
- The concept is mathematically **sound** and meaningful.

## WHY IS THAT A DIFFERENT POINT OF VIEW IN PHYSICS ?

- We do not need **points** and manifolds.
- We look for representations of **algebras** rather than **groups**.
- Symmetries are easily generalized in terms of **Hopf algebras**.
- Metric and gauge fields are unified in an elegant way.
- The theory is not abstract (not **formal** manipulations).
- The concept is mathematically **sound** and meaningful.
- Develops on **K-theory and K-cohomology**, Hochschild and cyclic homology, index theorems, pseudodifferential operators,  $C^*$ -algebras,

# SPECTRAL GEOMETRY

## THE SPECTRAL TRIPLE

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint unbounded operator  $D$ , satisfying several conditions:

- $\forall a \in \mathcal{A} [D, \pi(a)] \in B(\mathcal{H})$ ,  $D^{-1}$  is compact

# SPECTRAL GEOMETRY

## THE SPECTRAL TRIPLE

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint unbounded operator  $D$ , satisfying several conditions:

- 1  $\forall a \in \mathcal{A} [D, \pi(a)] \in B(\mathcal{H})$ ,  $D^{-1}$  is compact
- 2 even ST:  $\exists \gamma \in \mathcal{A}' : \gamma^2 = 1, \gamma = \gamma^\dagger, \gamma D + D\gamma = 0$ ,

# SPECTRAL GEOMETRY

## THE SPECTRAL TRIPLE

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint unbounded operator  $D$ , satisfying several conditions:

- 1  $\forall a \in \mathcal{A} [D, \pi(a)] \in B(\mathcal{H}), D^{-1}$  is compact
- 2 even ST:  $\exists \gamma \in \mathcal{A}' : \gamma^2 = 1, \gamma = \gamma^\dagger, \gamma D + D\gamma = 0,$
- 3  $\exists J$ , antilinear  $J^2 = \pm 1, JJ^\dagger = 1$   
 $J\gamma = \pm \gamma J, JD = \pm DJ, [J\pi(a)J, \pi(b)] = 0,$

# SPECTRAL GEOMETRY

## THE SPECTRAL TRIPLE

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint unbounded operator  $D$ , satisfying several conditions:

- 1  $\forall a \in \mathcal{A} [D, \pi(a)] \in B(\mathcal{H})$ ,  $D^{-1}$  is compact
- 2 even ST:  $\exists \gamma \in \mathcal{A}' : \gamma^2 = 1, \gamma = \gamma^\dagger, \gamma D + D\gamma = 0$ ,
- 3  $\exists J$ , antilinear  $J^2 = \pm 1, JJ^\dagger = 1$   
 $J\gamma = \pm\gamma J, JD = \pm DJ, [J\pi(a)J, \pi(b)] = 0$ ,
- 4  $[[D, a], J\pi(b)J] = 0$  ( $D$ : first order differential operator)

# SPECTRAL GEOMETRY

## THE SPECTRAL TRIPLE

Algebra  $\mathcal{A}$ , its faithful representation  $\pi$  on a Hilbert space  $\mathcal{H}$ , a selfadjoint unbounded operator  $D$ , satisfying several conditions:

- 1  $\forall a \in \mathcal{A} [D, \pi(a)] \in B(\mathcal{H})$ ,  $D^{-1}$  is compact
- 2 even ST:  $\exists \gamma \in \mathcal{A}' : \gamma^2 = 1, \gamma = \gamma^\dagger, \gamma D + D\gamma = 0$ ,
- 3  $\exists J$ , antilinear  $J^2 = \pm 1, JJ^\dagger = 1$   
 $J\gamma = \pm \gamma J, JD = \pm DJ, [J\pi(a)J, \pi(b)] = 0$ ,
- 4  $[[D, a], J\pi(b)J] = 0$  ( $D$ : first order differential operator)
- 5 ...+ conditions of „analysis” type

## THEOREM

If  $\mathcal{A} = C^\infty(M)$ ,  $M$  a spin Riemannian compact manifold,  $\mathcal{H} = L^2(S)$  (sections of spinor bundle) and  $D$  the Dirac operator on  $M$  then to  $(\mathcal{A}, \mathcal{H}, D)$  is a spectral triple (with a real structure).

# TWISTED SPECTRAL GEOMETRY

## DEFINITION (TWISTED REAL SPECTRAL TRIPLE)

Let  $\mathcal{A}$  be a  $*$ -algebra,  $(H, \pi)$  a representation of  $\mathcal{A}$ ,  $D$  a linear operator on  $H$ , and let  $\nu$  be a linear automorphism of  $H$ . We say that the triple  $(\mathcal{A}, H, D)$  admits a  $\nu$ -twisted real structure if there exists an anti-linear map  $J : H \rightarrow H$  such that  $J^2 = \epsilon \text{id}$ , and, for all  $a, b \in \mathcal{A}$ ,

# TWISTED SPECTRAL GEOMETRY

## DEFINITION (TWISTED REAL SPECTRAL TRIPLE)

Let  $\mathcal{A}$  be a  $*$ -algebra,  $(H, \pi)$  a representation of  $\mathcal{A}$ ,  $D$  a linear operator on  $H$ , and let  $\nu$  be a linear automorphism of  $H$ . We say that the triple  $(\mathcal{A}, H, D)$  admits a  $\nu$ -twisted real structure if there exists an anti-linear map  $J : H \rightarrow H$  such that  $J^2 = \epsilon \text{id}$ , and, for all  $a, b \in \mathcal{A}$ ,

$$[\pi(a), J\pi(b)J^{-1}] = 0,$$

# TWISTED SPECTRAL GEOMETRY

## DEFINITION (TWISTED REAL SPECTRAL TRIPLE)

Let  $\mathcal{A}$  be a  $*$ -algebra,  $(H, \pi)$  a representation of  $\mathcal{A}$ ,  $D$  a linear operator on  $H$ , and let  $\nu$  be a linear automorphism of  $H$ . We say that the triple  $(\mathcal{A}, H, D)$  admits a  $\nu$ -twisted real structure if there exists an anti-linear map  $J : H \rightarrow H$  such that  $J^2 = \epsilon \text{id}$ , and, for all  $a, b \in \mathcal{A}$ ,

$$[\pi(a), J\pi(b)J^{-1}] = 0,$$

$$[D, \pi(a)]J\nu^2(\pi(b))J^{-1} = J\pi(b)J^{-1}[D, \pi(a)],$$

# TWISTED SPECTRAL GEOMETRY

## DEFINITION (TWISTED REAL SPECTRAL TRIPLE)

Let  $\mathcal{A}$  be a  $*$ -algebra,  $(H, \pi)$  a representation of  $\mathcal{A}$ ,  $D$  a linear operator on  $H$ , and let  $\nu$  be a linear automorphism of  $H$ . We say that the triple  $(\mathcal{A}, H, D)$  admits a  $\nu$ -twisted real structure if there exists an anti-linear map  $J : H \rightarrow H$  such that  $J^2 = \epsilon \text{id}$ , and, for all  $a, b \in \mathcal{A}$ ,

$$[\pi(a), J\pi(b)J^{-1}] = 0,$$

$$[D, \pi(a)]J\nu^2(\pi(b))J^{-1} = J\pi(b)J^{-1}[D, \pi(a)],$$

$$DJ\nu = \epsilon'\nu JD,$$

where  $\epsilon, \epsilon' \in \{+, -\}$ .

# TWISTED SPECTRAL GEOMETRY

## DEFINITION (TWISTED REAL SPECTRAL TRIPLE)

Let  $\mathcal{A}$  be a  $*$ -algebra,  $(H, \pi)$  a representation of  $\mathcal{A}$ ,  $D$  a linear operator on  $H$ , and let  $\nu$  be a linear automorphism of  $H$ . We say that the triple  $(\mathcal{A}, H, D)$  admits a  $\nu$ -twisted real structure if there exists an anti-linear map  $J : H \rightarrow H$  such that  $J^2 = \epsilon \text{id}$ , and, for all  $a, b \in \mathcal{A}$ ,

$$[\pi(a), J\pi(b)J^{-1}] = 0,$$

$$[D, \pi(a)]J\nu^2(\pi(b))J^{-1} = J\pi(b)J^{-1}[D, \pi(a)],$$

$$DJ\nu = \epsilon'\nu JD,$$

where  $\epsilon, \epsilon' \in \{+, -\}$ .

$$\nu J\nu = J,$$

[Brzezinski, Dabrowski, Ciccoli, Sitarz, January 2016]

# TWISTED SPECTRAL GEOMETRY

Let  $\Omega_D^1$  be a bimodule of one forms:

$$\Omega_D^1 := \left\{ \sum_i \pi(a_i) [D, \pi(b_i)] \mid a_i, b_i \in A \right\}.$$

The standard fluctuation of a spectral triple  $(A, H, D)$  consist of adding to the Dirac operator  $D$  a selfadjoint one form  $\alpha \in \Omega_D^1$ .

# TWISTED SPECTRAL GEOMETRY

Let  $\Omega_D^1$  be a bimodule of one forms:

$$\Omega_D^1 := \left\{ \sum_i \pi(a_i)[D, \pi(b_i)] \mid a_i, b_i \in A \right\}.$$

The standard fluctuation of a spectral triple  $(A, H, D)$  consist of adding to the Dirac operator  $D$  a selfadjoint one form  $\alpha \in \Omega_D^1$ .

In case of a real spectral triple the fluctuated  $D$  is  $D + \alpha + \epsilon' J\alpha J^{-1}$ , where  $\alpha + \epsilon' J\alpha J^{-1}$  is selfadjoint.

# TWISTED SPECTRAL GEOMETRY

Let  $\Omega_D^1$  be a bimodule of one forms:

$$\Omega_D^1 := \left\{ \sum_i \pi(a_i)[D, \pi(b_i)] \mid a_i, b_i \in A \right\}.$$

The standard fluctuation of a spectral triple  $(A, H, D)$  consist of adding to the Dirac operator  $D$  a selfadjoint one form  $\alpha \in \Omega_D^1$ .

In case of a real spectral triple the fluctuated  $D$  is  $D + \alpha + \epsilon' J \alpha J^{-1}$ , where  $\alpha + \epsilon' J \alpha J^{-1}$  is selfadjoint.

For our case of  $\nu$ -twisted real spectral triple we set the fluctuated Dirac operator  $D_\alpha$  to be:

$$D_\alpha := D + \alpha + \epsilon' \nu J \alpha J^{-1} \nu,$$

with the requirement that  $\alpha + \epsilon' \nu J \alpha J^{-1} \nu$  is selfadjoint.

# THE TOOLS OF SPECTRAL GEOMETRY

## THE ACTION: HEAT TRACE

Take the Dirac operator  $D_A$  (that depends on physical degrees of freedom: **gauge fields** and **metric**:  $A$ ). Compute:

$$\lim_{t \rightarrow 0} \text{Tr} e^{-t(D_A)^2} = t^{-\frac{d}{2}} a_0(A) + t^{-\frac{d}{2}+1} a_2(A) + \dots$$

# THE TOOLS OF SPECTRAL GEOMETRY

## THE ACTION: HEAT TRACE

Take the Dirac operator  $D_A$  (that depends on physical degrees of freedom: **gauge fields** and **metric**:  $A$ ). Compute:

$$\lim_{t \rightarrow 0} \text{Tr} e^{-t(D_A)^2} = t^{-\frac{d}{2}} a_0(A) + t^{-\frac{d}{2}+1} a_2(A) + \dots$$

## THE ACTION: RESIDUES





Define for any number  $s$ , any operator  $T$  from the algebra  $\mathcal{A}$ :

$$\int_s T := \text{Res}_{z=s} \text{Tr} T |D_A|^{-z},$$

Then propose for the action

$$S(D_A) = \Lambda^d \int_d 1 + \Lambda^{d-2} \int_d (D_A)^2 + \Lambda^{d-4} \int_d (D_A)^4 + \dots$$

# THE SPECTRA



# GRAVITY

## THE ACTION FUNCTIONAL

For the geometries of the type  $M \times F$ , where  $M$  is a Riemannian manifold and  $F$  is a discrete geometry we obtain

# GRAVITY

## THE ACTION FUNCTIONAL

For the geometries of the type  $M \times F$ , where  $M$  is a Riemannian manifold and  $F$  is a discrete geometry we obtain

$$\begin{aligned} S &= \frac{1}{\pi^2} (48 f_4 \Lambda^4 - f_2 \Lambda^2 c + \frac{f_0}{4} d) \int \sqrt{g} d^4 x \\ &+ \frac{96 f_2 \Lambda^2 - f_0 c}{24 \pi^2} \int R \sqrt{g} d^4 x \\ &+ \frac{f_0}{10 \pi^2} \int \left( \frac{11}{6} R^* R^* - 3 C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma} \right) \sqrt{g} d^4 x + \dots \end{aligned}$$

# GRAVITY

## THE ACTION FUNCTIONAL

For the geometries of the type  $M \times F$ , where  $M$  is a Riemannian manifold and  $F$  is a discrete geometry we obtain

$$\begin{aligned} S &= \frac{1}{\pi^2} (48 f_4 \Lambda^4 - f_2 \Lambda^2 c + \frac{f_0}{4} d) \int \sqrt{g} d^4 x \\ &+ \frac{96 f_2 \Lambda^2 - f_0 c}{24 \pi^2} \int R \sqrt{g} d^4 x \\ &+ \frac{f_0}{10 \pi^2} \int (\frac{11}{6} R^* R^* - 3 C_{\mu\nu\rho\sigma} C^{\mu\nu\rho\sigma}) \sqrt{g} d^4 x + \dots \\ \dots &+ \frac{(-2 a f_2 \Lambda^2 + e f_0)}{\pi^2} \int |\varphi|^2 \sqrt{g} d^4 x + \frac{f_0}{2 \pi^2} \int a |D_\mu \varphi|^2 \sqrt{g} d^4 x \\ &- \frac{f_0}{12 \pi^2} \int a R |\varphi|^2 \sqrt{g} d^4 x + \frac{f_0}{2 \pi^2} \int b |\varphi|^4 \sqrt{g} d^4 x \\ &+ \frac{f_0}{2 \pi^2} \int (g_3^2 G_{\mu\nu}^i G^{\mu\nu i} + g_2^2 F_{\mu\nu}^\alpha F^{\mu\nu\alpha} + \frac{5}{3} g_1^2 B_{\mu\nu} B^{\mu\nu}) \sqrt{g} d^4 x \end{aligned}$$

## SPECTRAL ACTION FOR FLAT GEOMETRIES

There are **6 different** compact flat three-dimensional manifolds (**Bieberbach manifolds**). With P.Olczykowski we have demonstrated that the *symmetric* spectral action (depending on  $D^2$ ) does not distinguish which manifold we consider.

# GRAVITY




## SPECTRAL ACTION FOR FLAT GEOMETRIES

There are **6 different** compact flat three-dimensional manifolds (**Bieberbach manifolds**). With P.Olczykowski we have demonstrated that the *symmetric* spectral action (depending on  $D^2$ ) does not distinguish which manifold we consider.

## THE ASYMMETRIC ACTION AND $\eta$ INVARIANT

However, if the action depends on  $D$  (not only on even function of  $D$ ) then the difference is seen at the level of conformally invariant term – where a  $\eta$  invariant appears.

# THE SPECTRA



# SPECTRAL ACTION...

## ...AND THE NEUTRINO MASS

An interesting possibility occurs when one considers an extension of the spectral action for fermions, say:

$$S = \text{Tr} f \left( \frac{(D + P_\Psi)^2}{\Lambda^2} \right),$$

and then there are some interesting terms, which appear in the order  $\Lambda^{-2}$ .

# SPECTRAL ACTION...

## ...AND THE NEUTRINO MASS

An interesting possibility occurs when one considers an extension of the spectral action for fermions, say:

$$S = \text{Tr} f \left( \frac{(D + P_\Psi)^2}{\Lambda^2} \right),$$

and then there are some interesting terms, which appear in the order  $\Lambda^{-2}$ .

## WILCZEK TERMS

$$\mathcal{L}_m = \kappa (\bar{e}_L^c H^+ - \bar{\nu}^c H^0) (H^+ e_L - H^0 \nu),$$

where  $e_L, \nu$  is the doublet of left-handed leptons,  $H^+, H^0$  are the Higgs field components and  $\kappa$  is a coefficient (or a matrix if we take into account flavors)

# SPECTRAL ACTION

SPECULATE: (A.S: EUROP.PHYS.LETT., 86 (2009) 10007)

The speculation was such terms arise from quantum gravity corrections [Weinberg] gives the neutrino mass of the range of  $10^{-5}eV$ , which was much less than the experimental estimations. If we take the value of the cutoff parameter  $\Lambda = 10^{15}GeV$  and estimate the resulting neutrino mass (taking the coefficient in the term to be of order 1) we obtain the values of order  $10^{-2}eV$ , which agrees with the current experimental data.

# SPECTRAL ACTION

SPECULATE: (A.S: EUROP.PHYS.LETT., 86 (2009) 10007)

The speculation was such terms arise from quantum gravity corrections [Weinberg] gives the neutrino mass of the range of  $10^{-5}eV$ , which was much less than the experimental estimations. If we take the value of the cutoff parameter  $\Lambda = 10^{15}GeV$  and estimate the resulting neutrino mass (taking the coefficient in the term to be of order 1) we obtain the values of order  $10^{-2}eV$ , which agrees with the current experimental data.

## ...AND THE NEUTRINO MASS

Suppose we take it seriously - then it means that the nature of neutrino masses is completely different than the masses of fermions. Such terms do not affect the other fermions and the interactions (as these are corrections). However, they do affect the renormalizability of the theory.

# SPECTRAL ACTION

## ...AND THE TORSION:

The **usual** Dirac operator comes from the torsion-free connection. However, the operators with torsion are **in fact** almost indistinguishable from the **torsion-free** operators. In  $d = 3$  torsion is nothing but a scalar perturbation.

# SPECTRAL ACTION

## ...AND THE TORSION:

The **usual** Dirac operator comes from the torsion-free connection. However, the operators with torsion are **in fact** almost indistinguishable from the **torsion-free** operators. In  $d = 3$  torsion is nothing but a scalar perturbation.

LEMMA (A.SITARZ, A.ZAJĄC: LETT.MATH.PHYS. 98,3, 2011)

*The heat kernel coefficients for scalar perturbation of Dirac in dimension  $d = 3$  read:*

$$[a_1] = 2(4\pi)^{-\frac{3}{2}} \left( -\frac{1}{12}R + 2\phi^2 \right), \quad [a_2] = (4\pi)^{-\frac{3}{2}} \frac{8}{3} (\nabla_i \phi)(\nabla^i \phi).$$

# SPECTRAL ACTION

## ...AND THE TORSION:

The **usual** Dirac operator comes from the torsion-free connection. However, the operators with torsion are **in fact** almost indistinguishable from the **torsion-free** operators. In  $d = 3$  torsion is nothing but a scalar perturbation.

LEMMA (A.SITARZ, A.ZAJĄC: LETT.MATH.PHYS. 98,3, 2011)

*The heat kernel coefficients for scalar perturbation of Dirac in dimension  $d = 3$  read:*

$$[a_1] = 2(4\pi)^{-\frac{3}{2}} \left( -\frac{1}{12}R + 2\phi^2 \right), \quad [a_2] = (4\pi)^{-\frac{3}{2}} \frac{8}{3} (\nabla_i \phi)(\nabla^i \phi).$$

A. SITARZ, JPDO, 5,3, PP 305-317 (2014)

In  $d = 4$ , the noncommutative Wodzicki residue on the noncommutative torus, the action cannot be minimized.

# SPECTRAL ACTION

An interesting (though very special) example is that of the Standard Quantum Podleś Sphere [M. Eckstein, B. Iochum, A. Sitarz, CMP 332, 627– 668 \(2014\)](#)

# SPECTRAL ACTION

An interesting (though very special) example is that of the Standard Quantum Podleś Sphere [M. Eckstein, B. Iochum, A. Sitarz, CMP 332, 627– 668 \(2014\)](#)

## THE DIMENSION SPECTRUM

The dimension spectrum of the Podleś sphere is

$$\mathcal{S}_d(S_q^2) = -\mathbb{N} + i \frac{2\pi}{\log q} \mathbb{Z}$$

and the poles are at most of the second order.

# SPECTRAL ACTION

An interesting (though very special) example is that of the Standard Quantum Podleś Sphere [M. Eckstein, B. Iochum, A. Sitarz, CMP 332, 627– 668 \(2014\)](#)

## THE DIMENSION SPECTRUM

The dimension spectrum of the Podleś sphere is

$$\mathcal{S}_d(S_q^2) = -\mathbb{N} + i \frac{2\pi}{\log q} \mathbb{Z}$$

and the poles are at most of the second order.

## THE ACTION

$$\mathrm{Tr} e^{-t|\mathcal{D}_S|} = \frac{1}{\log^2 q} \left[ 2 \log^2(ut) + h_S(\log(ut)) \log(ut) + c_S(\log(ut)) \right] + \dots$$

where  $u = \frac{|w|q}{1-q^2}$  and  $h_S, c_S$  are periodic bounded  $C^\infty$ -functions on  $\mathbb{R}$

# SPECTRAL ACTION

## ... AND THE GAUSS-BONNET FOR THE NONCOMMUTATIVE TORI

Imagine we take a noncommutative torus ( $UV = \lambda VU$ ) algebra and take the following Dirac operator:

$$D_k = \sigma^1 \delta_1 + \sigma^2 \left( k \delta_2 + \frac{1}{2} \delta_2(k) \right)$$

# SPECTRAL ACTION

## ... AND THE GAUSS-BONNET FOR THE NONCOMMUTATIVE TORI

Imagine we take a noncommutative torus ( $UV = \lambda VU$ ) algebra and take the following Dirac operator:

$$D_k = \sigma^1 \delta_1 + \sigma^2 \left( k \delta_2 + \frac{1}{2} \delta_2(k) \right)$$

THEOREM (L.DABROWSKI, A.SITARZ, SIGMA 11 (2015), 075)

*The dressed scalar curvature for the asymmetric torus is:*

$$\tilde{R} = F_{11}(\Delta^{(1)}, \Delta^{(1)} \Delta^{(2)})(\delta_1(k) \cdot \delta_1(k)) + F_{22}(\Delta^{(1)}, \Delta^{(1)} \Delta^{(2)})(\delta_2(k), \delta_2(k)) + F'_{11}(\Delta^{(1)})(\delta_1(k)^2) + F'_{22}(\Delta^{(1)})(\delta_2(k)^2) + F_1(\Delta^{(1)})(\delta_{11}(k)) + F_2(\Delta^{(1)})(\delta_{22}(k)),$$

$$F_{11}(s, t) = -\frac{2\pi}{3k^3} \frac{(2s^2 + 4st + 4s + 3 + 8t + 3t^2)}{(t+1)^3(s+1)(s+t)}, \quad F_{22}(s, t) = \frac{\pi}{2k} \frac{(t^2 - 6t + 1)}{(t+1)^3},$$

$$F'_{11}(s) = \frac{4\pi}{3k^3} \frac{1}{(s+1)^3}, \quad F'_{22}(s) = -\frac{\pi}{2k} \frac{(s^2 - 6s + 1)}{(s+1)^3},$$

and

$$F_1(s) = \frac{2\pi}{3k^2} \frac{1}{(s+1)^2}, \quad F_2(s) = 0.$$

The trace of  $R$  vanishes.

# IS IT POINTLESS ?

- Geometry is **more** than what we have learned

# IS IT POINTLESS ?

- Geometry is **more** than what we have learned
- The input from physics **matters** - and tells us **what** geometry is

# IS IT POINTLESS ?

- Geometry is **more** than what we have learned
- The input from physics **matters** - and tells us **what** geometry is
- Physics is **not** the geometry we **imagine** it **should be**

# IS IT POINTLESS ?

- Geometry is **more** than what we have learned
- The input from physics **matters** - and tells us **what** geometry is
- Physics is **not** the geometry we **imagine** it **should be**
- Physics (as we observe) requires **new geometry**.

# IS IT POINTLESS ?


- Geometry is **more** than what we have learned
- The input from physics **matters** - and tells us **what** geometry is
- Physics is **not** the geometry we **imagine** it **should be**
- Physics (as we observe) requires **new geometry**.
- Reevaluate more possibilities: Einstein -Cartan

# IS IT POINTLESS ?

- Geometry is **more** than what we have learned
- The input from physics **matters** - and tells us **what** geometry is
- Physics is **not** the geometry we **imagine** it **should be**
- Physics (as we observe) requires **new geometry**.
- Reevaluate more possibilities: Einstein -Cartan
- Benefits (or drawbacks): **NO supersymmetry (!)**

# IS IT POINTLESS ?

- Geometry is **more** than what we have learned
- The input from physics **matters** - and tells us **what** geometry is
- Physics is **not** the geometry we **imagine** it **should be**
- Physics (as we observe) requires **new geometry**.
- Reevaluate more possibilities: Einstein -Cartan
- Benefits (or drawbacks): **NO supersymmetry (!)**
- Future: **Geometry as Quantum Field Theory**



THANK YOU